

Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis (Review)

Brosseau L, Welch V, Wells GA, de Bie R, Gam A, Harman K, Morin M, Shea B, Tugwell P



**THE COCHRANE
COLLABORATION®**

This is a reprint of a Cochrane review, prepared and maintained by The Cochrane Collaboration and published in *The Cochrane Library* 2009, Issue 1

<http://www.thecochranelibrary.com>



Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis (Review)
Copyright © 2009 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

TABLE OF CONTENTS

HEADER	1
ABSTRACT	1
PLAIN LANGUAGE SUMMARY	2
BACKGROUND	4
OBJECTIVES	4
METHODS	4
RESULTS	5
DISCUSSION	7
AUTHORS' CONCLUSIONS	8
ACKNOWLEDGEMENTS	8
REFERENCES	9
CHARACTERISTICS OF STUDIES	11
DATA AND ANALYSES	19
Analysis 1.1. Comparison 1 Laser vs Placebo- End of treatment (approx 10 wks), Outcome 1 Pain (Visual Analogue Scale 10 cm).	25
Analysis 1.2. Comparison 1 Laser vs Placebo- End of treatment (approx 10 wks), Outcome 2 Pain (0-12 scale).	26
Analysis 1.3. Comparison 1 Laser vs Placebo- End of treatment (approx 10 wks), Outcome 3 McGill Pain questionnaire.	26
Analysis 1.4. Comparison 1 Laser vs Placebo- End of treatment (approx 10 wks), Outcome 4 Ritchie Index.	27
Analysis 1.5. Comparison 1 Laser vs Placebo- End of treatment (approx 10 wks), Outcome 5 Health Assessment Questionnaire (HAQ).	27
Analysis 1.6. Comparison 1 Laser vs Placebo- End of treatment (approx 10 wks), Outcome 6 MCP ROM (degrees).	28
Analysis 1.7. Comparison 1 Laser vs Placebo- End of treatment (approx 10 wks), Outcome 7 PIP ROM (degrees).	28
Analysis 1.8. Comparison 1 Laser vs Placebo- End of treatment (approx 10 wks), Outcome 8 Knee ROM.	29
Analysis 1.9. Comparison 1 Laser vs Placebo- End of treatment (approx 10 wks), Outcome 9 Ankle ROM.	30
Analysis 1.10. Comparison 1 Laser vs Placebo- End of treatment (approx 10 wks), Outcome 10 Flexibility- tip to palm distance (cm).	32
Analysis 1.11. Comparison 1 Laser vs Placebo- End of treatment (approx 10 wks), Outcome 11 Morning stiffness duration (min).	32
Analysis 1.12. Comparison 1 Laser vs Placebo- End of treatment (approx 10 wks), Outcome 12 Morning stiffness not improved.	33
Analysis 1.13. Comparison 1 Laser vs Placebo- End of treatment (approx 10 wks), Outcome 13 Rheumatoid factor positive.	33
Analysis 1.14. Comparison 1 Laser vs Placebo- End of treatment (approx 10 wks), Outcome 14 Grip strength.	34
Analysis 1.15. Comparison 1 Laser vs Placebo- End of treatment (approx 10 wks), Outcome 15 Suprapatellar swelling - right knee (cm).	35
Analysis 1.16. Comparison 1 Laser vs Placebo- End of treatment (approx 10 wks), Outcome 16 Suprapatellar swelling- left knee (cm).	36
Analysis 1.17. Comparison 1 Laser vs Placebo- End of treatment (approx 10 wks), Outcome 17 MCP swelling (cm).	36
Analysis 1.18. Comparison 1 Laser vs Placebo- End of treatment (approx 10 wks), Outcome 18 PIP swelling (cm).	37
Analysis 1.19. Comparison 1 Laser vs Placebo- End of treatment (approx 10 wks), Outcome 19 Walking speed (sec).	37
Analysis 1.20. Comparison 1 Laser vs Placebo- End of treatment (approx 10 wks), Outcome 20 Fibrinogen.	38
Analysis 1.21. Comparison 1 Laser vs Placebo- End of treatment (approx 10 wks), Outcome 21 Leukocytes.	38
Analysis 1.22. Comparison 1 Laser vs Placebo- End of treatment (approx 10 wks), Outcome 22 Lymphocytes.	39
Analysis 1.23. Comparison 1 Laser vs Placebo- End of treatment (approx 10 wks), Outcome 23 ESR (mm/hr).	39
Analysis 1.24. Comparison 1 Laser vs Placebo- End of treatment (approx 10 wks), Outcome 24 CRP (g/mL).	40
Analysis 1.25. Comparison 1 Laser vs Placebo- End of treatment (approx 10 wks), Outcome 25 Hemoglobin.	40
Analysis 1.26. Comparison 1 Laser vs Placebo- End of treatment (approx 10 wks), Outcome 26 Platelets.	41
Analysis 2.1. Comparison 2 Laser vs Placebo- End of follow-up (approx 20 wks), Outcome 1 Pain.	41
Analysis 2.2. Comparison 2 Laser vs Placebo- End of follow-up (approx 20 wks), Outcome 2 McGill Pain questionnaire.	42
Analysis 2.3. Comparison 2 Laser vs Placebo- End of follow-up (approx 20 wks), Outcome 3 Health Assessment Questionnaire (HAQ).	43

Analysis 2.4. Comparison 2 Laser vs Placebo- End of follow-up (approx 20 wks), Outcome 4 Ritchie Index.	43
Analysis 2.5. Comparison 2 Laser vs Placebo- End of follow-up (approx 20 wks), Outcome 5 PIP ROM (degrees).	44
Analysis 2.6. Comparison 2 Laser vs Placebo- End of follow-up (approx 20 wks), Outcome 6 Knee ROM.	44
Analysis 2.7. Comparison 2 Laser vs Placebo- End of follow-up (approx 20 wks), Outcome 7 Ankle ROM.	46
Analysis 2.8. Comparison 2 Laser vs Placebo- End of follow-up (approx 20 wks), Outcome 8 MCP ROM (degrees).	47
Analysis 2.9. Comparison 2 Laser vs Placebo- End of follow-up (approx 20 wks), Outcome 9 Morning stiffness duration (min).	48
Analysis 2.10. Comparison 2 Laser vs Placebo- End of follow-up (approx 20 wks), Outcome 10 Walking speed (sec).	48
Analysis 2.11. Comparison 2 Laser vs Placebo- End of follow-up (approx 20 wks), Outcome 11 Grip strength.	49
Analysis 2.12. Comparison 2 Laser vs Placebo- End of follow-up (approx 20 wks), Outcome 12 Suprapatellar swelling- left knee (cm).	50
Analysis 2.13. Comparison 2 Laser vs Placebo- End of follow-up (approx 20 wks), Outcome 13 Suprapatellar swelling - right knee (cm).	50
Analysis 2.14. Comparison 2 Laser vs Placebo- End of follow-up (approx 20 wks), Outcome 14 MCP swelling (cm).	51
Analysis 2.15. Comparison 2 Laser vs Placebo- End of follow-up (approx 20 wks), Outcome 15 PIP swelling (cm).	51
Analysis 2.16. Comparison 2 Laser vs Placebo- End of follow-up (approx 20 wks), Outcome 16 Thermographic index.	52
Analysis 2.17. Comparison 2 Laser vs Placebo- End of follow-up (approx 20 wks), Outcome 17 Rheumatoid factor positive.	52
Analysis 2.18. Comparison 2 Laser vs Placebo- End of follow-up (approx 20 wks), Outcome 18 ESR (mm/hr).	53
Analysis 2.19. Comparison 2 Laser vs Placebo- End of follow-up (approx 20 wks), Outcome 19 CRP (g/mL).	53
Analysis 2.20. Comparison 2 Laser vs Placebo- End of follow-up (approx 20 wks), Outcome 20 Hemoglobin.	54
Analysis 2.21. Comparison 2 Laser vs Placebo- End of follow-up (approx 20 wks), Outcome 21 Platelets.	54
Analysis 3.1. Comparison 3 Treatment length- subgroup analysis for end of treatment results, Outcome 1 Pain (10 cm VAS or 0-12 scale).	55
Analysis 4.1. Comparison 4 Treatment length- subgroup analysis for end of follow up results, Outcome 1 Pain (10 cm VAS or 0-12 scale).	56
Analysis 5.1. Comparison 5 Methodologic quality: low (<3) vs high (>= 3), Outcome 1 Pain at end of treatment.	57
Analysis 6.1. Comparison 6 Laser vs Placebo - Joint vs Nerve - end of treatment, Outcome 1 Pain (10 cm VAS or 0-12 scale).	59
Analysis 7.1. Comparison 7 Laser vs Placebo - Joint vs Nerve - end of follow up, Outcome 1 Pain (10 cm VAS or 0-12 scale).	61
Analysis 8.1. Comparison 8 Laser vs Placebo- Wavelength analysis - End of treatment, Outcome 1 Pain.	63
Analysis 8.2. Comparison 8 Laser vs Placebo- Wavelength analysis - End of treatment, Outcome 2 McGill Pain questionnaire.	65
Analysis 8.3. Comparison 8 Laser vs Placebo- Wavelength analysis - End of treatment, Outcome 3 Ritchie Index.	66
Analysis 8.4. Comparison 8 Laser vs Placebo- Wavelength analysis - End of treatment, Outcome 4 Health Assessment Questionnaire (HAQ).	67
Analysis 8.5. Comparison 8 Laser vs Placebo- Wavelength analysis - End of treatment, Outcome 5 MCP ROM (degrees).	68
Analysis 8.6. Comparison 8 Laser vs Placebo- Wavelength analysis - End of treatment, Outcome 6 PIP ROM (degrees).	69
Analysis 8.7. Comparison 8 Laser vs Placebo- Wavelength analysis - End of treatment, Outcome 7 Left knee Range of motion.	70
Analysis 8.8. Comparison 8 Laser vs Placebo- Wavelength analysis - End of treatment, Outcome 8 Right knee ROM.	71
Analysis 8.9. Comparison 8 Laser vs Placebo- Wavelength analysis - End of treatment, Outcome 9 Left ankle ROM.	72
Analysis 8.10. Comparison 8 Laser vs Placebo- Wavelength analysis - End of treatment, Outcome 10 Right ankle ROM.	73
Analysis 8.11. Comparison 8 Laser vs Placebo- Wavelength analysis - End of treatment, Outcome 11 Flexibility- tip to palm distance (cm).	74
Analysis 8.12. Comparison 8 Laser vs Placebo- Wavelength analysis - End of treatment, Outcome 12 Morning stiffness duration (min).	75
Analysis 8.13. Comparison 8 Laser vs Placebo- Wavelength analysis - End of treatment, Outcome 13 Morning stiffness not improved.	77
Analysis 8.14. Comparison 8 Laser vs Placebo- Wavelength analysis - End of treatment, Outcome 14 Rheumatoid factor positive.	78

Analysis 8.15. Comparison 8 Laser vs Placebo- Wavelength analysis - End of treatment, Outcome 15 Grip strength (mmHg, KPa, kg).	79
Analysis 8.16. Comparison 8 Laser vs Placebo- Wavelength analysis - End of treatment, Outcome 16 Suprapatellar swelling - right knee (cm).	81
Analysis 8.17. Comparison 8 Laser vs Placebo- Wavelength analysis - End of treatment, Outcome 17 Suprapatellar swelling- left knee (cm).	82
Analysis 8.18. Comparison 8 Laser vs Placebo- Wavelength analysis - End of treatment, Outcome 18 MCP swelling (cm).	83
Analysis 8.19. Comparison 8 Laser vs Placebo- Wavelength analysis - End of treatment, Outcome 19 PIP swelling (cm).	84
Analysis 8.20. Comparison 8 Laser vs Placebo- Wavelength analysis - End of treatment, Outcome 20 Walking speed (sec).	85
Analysis 8.21. Comparison 8 Laser vs Placebo- Wavelength analysis - End of treatment, Outcome 21 Fibrinogen.	86
Analysis 8.22. Comparison 8 Laser vs Placebo- Wavelength analysis - End of treatment, Outcome 22 Leukocytes.	87
Analysis 8.23. Comparison 8 Laser vs Placebo- Wavelength analysis - End of treatment, Outcome 23 Lymphocytes.	88
Analysis 8.24. Comparison 8 Laser vs Placebo- Wavelength analysis - End of treatment, Outcome 24 ESR (mm/hr).	89
Analysis 8.25. Comparison 8 Laser vs Placebo- Wavelength analysis - End of treatment, Outcome 25 CRP (g/mL).	90
Analysis 8.26. Comparison 8 Laser vs Placebo- Wavelength analysis - End of treatment, Outcome 26 Hemoglobin.	92
Analysis 8.27. Comparison 8 Laser vs Placebo- Wavelength analysis - End of treatment, Outcome 27 Platelets.	94
Analysis 9.1. Comparison 9 Subgroup Dosage Analysis, Outcome 1 Pain VAS <= 3 J/cm2.	95
Analysis 9.2. Comparison 9 Subgroup Dosage Analysis, Outcome 2 Pain VAS > 3 J/cm2.	95
Analysis 9.3. Comparison 9 Subgroup Dosage Analysis, Outcome 3 ESR low dose.	96
Analysis 9.4. Comparison 9 Subgroup Dosage Analysis, Outcome 4 ESR high dose.	96
Analysis 9.5. Comparison 9 Subgroup Dosage Analysis, Outcome 5 CRP low dose.	97
Analysis 9.6. Comparison 9 Subgroup Dosage Analysis, Outcome 6 CRP high dose.	97
Analysis 9.7. Comparison 9 Subgroup Dosage Analysis, Outcome 7 HAQ high dose.	98
Analysis 9.8. Comparison 9 Subgroup Dosage Analysis, Outcome 8 Hemoglobin high dose.	98
Analysis 9.9. Comparison 9 Subgroup Dosage Analysis, Outcome 9 Morning stiffness (hours) high dose.	99
Analysis 9.10. Comparison 9 Subgroup Dosage Analysis, Outcome 10 Grip strength (kg) low.	99
Analysis 9.11. Comparison 9 Subgroup Dosage Analysis, Outcome 11 Grip strength (kg) high.	100
Analysis 9.12. Comparison 9 Subgroup Dosage Analysis, Outcome 12 MCP swelling (cm) low.	100
Analysis 9.13. Comparison 9 Subgroup Dosage Analysis, Outcome 13 Flexibility tip to palm (cm) low.	101
Analysis 9.14. Comparison 9 Subgroup Dosage Analysis, Outcome 14 Flexibility tip to palm high dose.	101
Analysis 10.1. Comparison 10 Laser vs Placebo- contralateral control (End of treatment), Outcome 1 Number of patients improved in pain relief.	102
Analysis 10.2. Comparison 10 Laser vs Placebo- contralateral control (End of treatment), Outcome 2 Number of patients improved in morning stiffness.	102
APPENDICES	102
WHAT'S NEW	103
HISTORY	103
CONTRIBUTIONS OF AUTHORS	104
DECLARATIONS OF INTEREST	104
SOURCES OF SUPPORT	104
INDEX TERMS	104

[Intervention Review]

Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Lucie Brosseau¹, Vivian Welch², George A Wells³, Rob de Bie⁴, Arne Gam⁵, Katherine Harman⁶, Michelle Morin⁷, Beverley Shea⁸, Peter Tugwell²

¹School of Rehabilitation Sciences, Faculty of Health Sciences, University of Ottawa, Ottawa, Canada. ²Centre for Global Health, Institute of Population Health, University of Ottawa, Ottawa, Canada. ³Cardiovascular Research Reference Centre, University of Ottawa Heart Institute, Ottawa, Canada. ⁴Department of Epidemiology, Maastricht University, Maastricht, Netherlands. ⁵Vanlose, Denmark. ⁶Psychology, Neuroscience, Dalhousie University, Halifax, Canada. ⁷School of Rehabilitation Sciences, University of Ottawa, Ottawa, Canada. ⁸Institute of Population Health, University of Ottawa, Ottawa, Canada

Contact address: Lucie Brosseau, School of Rehabilitation Sciences, Faculty of Health Sciences, University of Ottawa, 451 Smyth Road, Ottawa, Ontario, K1H 8M5, Canada. lbrossea@uottawa.ca. (Editorial group: Cochrane Musculoskeletal Group.)

Cochrane Database of Systematic Reviews, Issue 1, 2009 (Status in this issue: *Edited*)

Copyright © 2009 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

DOI: 10.1002/14651858.CD002049.pub2

This version first published online: 19 October 2005 in Issue 4, 2005. Re-published online with edits: 21 January 2009 in Issue 1, 2009.

Last assessed as up-to-date: 21 August 2005. (Help document - [Dates and Statuses](#) explained)

This record should be cited as: Brosseau L, Welch V, Wells GA, de Bie R, Gam A, Harman K, Morin M, Shea B, Tugwell P. Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis. *Cochrane Database of Systematic Reviews* 2005, Issue 4. Art. No.: CD002049. DOI: 10.1002/14651858.CD002049.pub2.

ABSTRACT

Background

Rheumatoid arthritis (RA) affects a large proportion of the population. Low Level Laser Therapy (LLLT) was introduced as an alternative non-invasive treatment for RA about ten years ago. LLLT is a light source that generates extremely pure light, of a single wavelength. The effect is not thermal, but rather related to photochemical reactions in the cells. The effectiveness of LLLT for rheumatoid arthritis is still controversial. This review is an update of the original review published in October 1998.

Objectives

To assess the effectiveness of LLLT in the treatment of RA.

Search strategy

We initially searched MEDLINE, EMBASE (from 1998), the registries of the Cochrane Musculoskeletal Group and the field of Rehabilitation and Related Therapies as well as the Cochrane Central Register of Controlled Trials (CENTRAL) up to June 2001. This search has now been updated to include articles published up to June 2005.

Selection criteria

Following an a priori protocol, only randomized controlled trials of LLLT for the treatment of patients with a clinical diagnosis of RA were eligible. Abstracts were excluded unless further data could be obtained from the authors.

Data collection and analysis

Two reviewers independently selected trials for inclusion, then extracted data and assessed quality using predetermined forms. Heterogeneity was tested using chi-squared. A fixed effects model was used throughout for continuous variables, except where heterogeneity

existed, in which case, a random effects model was used. Results were analyzed as weighted mean differences (WMD) with 95% confidence intervals (CI), where the difference between the treated and control groups was weighted by the inverse of the variance. Dichotomous outcomes were analyzed with relative risks.

Main results

A total of 222 patients were included in the five placebo-controlled trials, with 130 randomized to laser therapy. Relative to a separate control group, LLLT reduced pain by 1.10 points (95% CI: 1.82, 0.39) on visual analogue scale relative to placebo, reduced morning stiffness duration by 27.5 minutes (95%CI: 2.9 to 52 minutes) and increased tip to palm flexibility by 1.3 cm (95% CI: 0.8 to 1.7). Other outcomes such as functional assessment, range of motion and local swelling did not differ between groups. There were no significant differences between subgroups based on LLLT dosage, wavelength, site of application or treatment length. For RA, relative to a control group using the opposite hand, there was no difference observed between the control and treatment hand for morning stiffness duration, and also no significant improvement in pain relief RR 13.00 (95% CI: 0.79 to 214.06). However, only one study was included as using the contralateral limb as control. .

Authors' conclusions

LLLT could be considered for short-term treatment for relief of pain and morning stiffness for RA patients, particularly since it has few side-effects. Clinicians and researchers should consistently report the characteristics of the LLLT device and the application techniques used. New trials on LLLT should make use of standardized, validated outcomes. Despite some positive findings, this meta-analysis lacked data on how LLLT effectiveness is affected by four important factors: wavelength, treatment duration of LLLT, dosage and site of application over nerves instead of joints. There is clearly a need to investigate the effects of these factors on LLLT effectiveness for RA in randomized controlled clinical trials.

PLAIN LANGUAGE SUMMARY

Low level laser therapy for rheumatoid arthritis

Does low level laser therapy work for treating rheumatoid arthritis?

Six studies of medium quality were reviewed and provide the best evidence we have today. Collectively, these studies tested over 220 people with rheumatoid arthritis. The studies compared how well people did while receiving either laser therapy or a 'placebo' (fake) laser therapy. Laser therapy was given mostly on the hands and generally for two to three times a week for four weeks. There were also many different wavelengths and dosages given.

What is rheumatoid arthritis and low level laser therapy?

Rheumatoid arthritis (RA) is a disease in which the body's immune system attacks its own healthy tissues. The attack happens mostly in the joints of the hands and feet and causes redness, pain, swelling and heat around the joints. Drug and non-drug treatments are used to relieve pain and/or swelling. Low level laser therapy, is a non-drug treatment used to decrease swelling and pain. Without producing heat, the laser emits very pure light that causes light and chemical reactions in cells where it is targeted.

What did the studies show?

Studies showed that laser therapy decreased pain and morning stiffness more than 'placebo' laser therapy. Laser therapy also increased hand flexibility more than placebo therapy.

Pain decreased by 1.10 points on a scale of 1-10. The length of time for morning stiffness decreased by 28 minutes.

Studies also showed that laser therapy worked just as well as 'placebo' laser therapy to improve range of motion, function, swelling and grip strength.

Only two of the studies measured the effect of laser therapy three months after the end of treatment. The results from these studies indicated that laser therapy worked just as well as 'placebo' therapy after three months times.

Dose, length of laser administration time and wavelength of the laser therapies did not appear to make a significant difference, though there was some evidence indicating that longer administration times and shorter wavelengths produced better effects.

Were there any side effects?

No side effects were reported in the studies.

What is the bottom line?

There is 'silver' level evidence that low level laser therapy in people with rheumatoid arthritis for up to four weeks does decrease pain and morning stiffness. It does not appear, however, to have long-lasting effects.

Most of the studies tested laser therapy on the hand, so it is not clear whether laser therapy would affect other joints of the body the same way.

BACKGROUND

Rheumatoid arthritis (RA) affects a large proportion of the population. The prevalence of RA is increasing with the aging population in industrial countries, especially with the baby boomers phenomenon (Walker 1996). Indeed, the number of Americans who will be diagnosed with arthritis will reach 59.4 million in 2020 compared to the 38 million diagnosed in 1990. This represents an increase of more than 59%. Thus, 18.2% of the population will be affected by arthritis: a proportion of one in five persons (Walker 1996). RA is recognized as an important source of disability and handicap which leads to considerable socioeconomic costs due to medical treatments, surgical interventions and frequent absences from work (Badley 1994).

A large number of pharmacological treatments are available for RA, ranging from analgesics to disease modifying anti-rheumatic drugs (DMARDs). The effectiveness and severity of adverse side effects of these interventions varies.

A new non invasive therapy was introduced ten years ago to existing physical agents used in medicine and physiotherapy in the treatment of RA: Low Level Laser Therapy (LLLT), Classes I, II and III (Brosseau 1994). Low Level Laser Therapy (LLLT) is a light source that generates extremely pure light, of a single wavelength. The effect is not thermal, but rather related to photochemical reactions in the cells. The effectiveness of laser therapy is still unclear (Beckerman 1992) and needs to be examined more rigorously.

In recent years, there has been increasing recognition that LLLT may have physiologic effects mediated by photochemical actions at the cellular level in animal and human tissues. Bassler et al. concluded that LLLT also has cartilage stimulatory properties in human beings (Bassler 1985). Some studies suggest that LLLT has a beneficial anti-inflammatory and pain attenuation effect in human beings (Ceccherelli 1989, King 1990, Tsurko 1983). Some studies in animals, conducted to elucidate the possible mechanism of action of LLLT irradiation, suggest also the alteration of nerve tissue (Kudoh 1989).

A meta-analysis on the effect of LLLT on musculoskeletal pain showed that LLLT has no effect on musculoskeletal pain syndromes (Gam 1993). However, this meta-analysis did not examine rheumatoid arthritis specifically. Additionally, trials which used the contralateral limb as a control were combined with those with a separate control group.

OBJECTIVES

The aim is to assess the effectiveness of Low Level Laser Therapy (LLLT) in the treatment of rheumatoid arthritis (RA).

A secondary purpose is to determine the most effective method of administering LLLT for RA, including the optimal:

- wavelength
- dosage
- application techniques
- length of treatment

METHODS

Criteria for considering studies for this review

Types of studies

Following an a priori protocol, studies were eligible if they were Randomized Controlled Trials (RCTs).

Types of participants

Only trials with subjects aged 18 years and over, with clinical or radiological confirmation of the diagnosis of RA were included. Trials published in languages other than French and English were not analyzed because of the time and cost involved in translation. Abstracts were excluded unless further data could be obtained from the authors.

Types of interventions

All types of LLLT (Classes I, II, and III) including all wavelengths from 632nm to 1064nm, were included in this review. Trials which compared one intervention group using LLLT with one standard treatment were included. Trials which were placebo-controlled were also included.

Trials which used the opposite limb as a control were analyzed separately since laser may have systemic effects (Prwemstay 1987, Young 1988).

Types of outcome measures

The primary endpoints for measurement of effectiveness were the outcome measures recommended by the conference on Outcome Measures for Rheumatoid Arthritis Clinical Trials 1993 (OMERACT 1993).

These measures include:

- a) Number of tender joints
- b) Number of swollen joints
- c) Pain
- d) Physician global assessment
- e) Patient global assessment
- f) Functional status (for example, measured by the Activities of Daily Living Scale)
- g) Acute phase reactants
- h) Radiological damage

In addition to these outcomes, one of the authors (LB) developed a theoretical framework of important physiotherapy outcomes for

the study of LLLT for the treatment of RA (Brosseau 1998). These outcomes were assessed as secondary endpoints and include:

- 1) Articular mobility
- 2) Muscular testing
- 3) Systemic components
- 4) Medication usage
- 5) Adverse side effects

Search methods for identification of studies

We initially searched MEDLINE up to December, 2001, EMBASE from 1988 to December 2001 and the Cochrane Central Register of Controlled Trials Register (CENTRAL), Issue 4, 1998. Details of the search strategy are given in the appendix. The electronic search was complemented by the following hand searches: 1) Bibliographic references; 2) Current Contents up to December 2001 (to identify articles not yet indexed in MEDLINE); 3) Abstracts published in special issues of specialized journals or in Conference Proceedings; 4) Coordinating offices of the trials registries of the Cochrane Field of Physical and Related Therapies and the Cochrane Musculoskeletal Group were also contacted. This exact search has now been updated to include articles published up until June 2005.

Reference lists were hand-searched for further identification of published work, presentations at scientific meetings and personal communications. Content experts were contacted for additional studies and unpublished data. If data could not be obtained, abstracts were not used.

The search strategy used is in [Appendix 1](#).

Data collection and analysis

The trials retrieved by the above search strategy were assessed independently by two reviewers (LB, RD) according to the selection criteria. Differences were resolved by consensus.

From each included trial, we collected information regarding the trial design, patient characteristics, dosages and treatment periods, and baseline and end of study outcomes. Data concerning details of the study population, intervention and outcomes were extracted independently by two reviewers (LB, VR). Differences in data extraction were resolved by referring back to the original article and establishing consensus. A third reviewer (BS) was consulted to help resolve differences.

Data on the outcomes from each trial were pooled to arrive at an overall estimate of the effectiveness of laser therapy. Where possible, the analyses were based on intention-to-treat data from the individual clinical trials.

Since laser is thought to have systemic effects, we decided a priori to analyze trials separately depending on whether the control group consisted of a separate group of patients or whether the control group consisted of the patients treated with active laser on one side and sham laser on the other side (contralateral control trials).

Subgroup analyses were attempted to determine the effects of the method of LLLT administration, methodological quality and length of treatment on outcomes.

For continuous data, results are presented as a weighted mean difference [WMD]. For dichotomous data, a relative risk (RR) was calculated (Mantel 1959). A test for heterogeneity was calculated using a Chi square test, with a significance level of $p < .05$. Fixed effects models were used throughout unless heterogeneity was significant, in which case, a random effects model was used. Publication bias was not assessed due to the small number of included studies.

RESULTS

Description of studies

See: [Characteristics of included studies](#); [Characteristics of excluded studies](#).

The initial literature search retrieved 191 articles. Of these, six trials met the inclusion criteria (Bliddal 1987, Goats 1996, Hall 1994, Johannsen 1994, Palmgren 1989, Walker 1987). Five were randomized controlled trials of laser therapy versus a placebo laser apparatus with a separate control group (Goats 1996, Hall 1994, Johannsen 1994, Palmgren 1989, Walker 1987). One used the contralateral joint as the control (Bliddal 1987). Five studies were excluded in the initial search: one study was an early abstract of one of the included studies (Walker 1983), another lacked a control group (Asada K 1989), two had insufficient statistical data (Goldman 1980; Heussler 1993) while the last one was not randomized as indicated by the title of the article (Oyamada 1988). The updated search to June 2005 identified an additional six potential articles which had to be excluded for the following reasons: three studies were literature reviews (Ammer K 2003, Day R 1999, De Bie 1998), two articles were only available in Russian (Sidorov VD 1999; Sidorov VD 2000), and the final article included patients with juvenile RA (Fulga C 1998). Moreover, in the initial review, there were also several studies waiting assessment due to a lack of information - either because they were abstracts (Agambar 1992, Matulis 1983), or because they were published in German or Japanese (Barabas 1988, Miyagi 1989, Taghawinejad 1985). Letters were sent out at that time to the respective authors to obtain further information. However, the authors did not respond, preventing any further updated assessment by the authors of this review.

Of the included trials, several reported outcomes as graphs. The mean scores and standard deviations were estimated from the graphs. One study presented the logarithm of pain measured on a 10 cm VAS (Walker 1987). We assumed the logarithm involved the natural exponent and translated these by performing an antilog operation. Another study reported pain on a VAS scale, but

then transformed these results to report only 3 categories: laser better than placebo, laser worse than placebo and no difference (Bliddal 1987). One study reported end of study scores corrected and uncorrected for disease activity (Johannsen 1994). Because randomization is assumed to control for disease activity and other variables, the uncorrected values were used for this meta-analysis. A total of 204 patients were included in the five placebo-controlled trials, with 112 randomized to laser therapy. For the trial which used the opposite limb as a control, 18 patients were included in the analysis. The mean age ranged from 53 to 67 years and baseline morning stiffness ranged from 60 to 90 minutes. The schedule of treatment was 2-3 sessions per week for 3-4 weeks for all trials except Walker et al 1987 which treated patients for 10 weeks, 3 times per week. All trials included patients with RA of the hands or thumb except Walker et al 1987, which included patients with a diagnosis of RA, but did not specify which joints were affected. The outcomes measured in the included studies were as follows: pain was reported by six trials, functional status by two, range of motion by four, swelling by three, grip strength by three, and morning stiffness by four.

Risk of bias in included studies

The quality of the included studies was assessed using a validated scale (Jadad 1996), which includes the appropriateness of randomization, appropriateness of blinding and description of dropouts and withdrawals. Quality was assessed independently by two reviewers (LB, VR). Differences were resolved by consensus. A third reviewer (BS) was consulted when necessary.

The median methodological quality was 3, with a range from 1-5 (Table 1). The individual components of the Jadad score of randomization, blinding and withdrawals are shown in Table 1. Only one trial scored full points for randomization. Five trials scored full points for double-blinding and four trials did not report withdrawals and dropouts. A subgroup analysis was conducted to determine whether the methodological quality influenced the effect of laser by dividing the studies into low and high quality at the median score.

To rank the strength of scientific evidence we used the grading system adopted by the Cochrane Musculoskeletal Group (Tugwell 2004). This grading system is based on four levels: Platinum, Gold, Silver and Bronze.

Platinum: A published systematic review that has at least two individual controlled trials each satisfying the following :

Sample sizes of at least 50 per group - if these do not find a statistically significant difference, they are adequately powered for a 20% relative difference in the relevant outcome.

Blinding of patients and assessors for outcomes.

Handling of withdrawals >80% follow up (imputations based on methods such as Last Observation Carried Forward (LOCF) are acceptable).

Concealment of treatment allocation.

Gold: At least one randomised clinical trial meeting all of the following criteria for the major outcome(s) as reported:

Sample sizes of at least 50 per group - if these do not find a statistically significant difference, they are adequately powered for a 20% relative difference in the relevant outcome.

Blinding of patients and assessors for outcomes.

Handling of withdrawals > 80% follow up (imputations based on methods such as LOCF are acceptable).

Concealment of treatment allocation.

Silver: A systematic review or randomised trial that does not meet the above criteria. Silver ranking would also include evidence from at least one study of non-randomised cohorts that did and did not receive the therapy, or evidence from at least one high quality case-control study. A randomised trial with a 'head-to-head' comparison of agents would be considered silver level ranking unless a reference were provided to a comparison of one of the agents to placebo showing at least a 20% relative difference.

Bronze: The bronze ranking is given to evidence if at least one high quality case series without controls (including simple before/after studies in which patients act as their own control) or if the conclusion is derived from expert opinion based on clinical experience without reference to any of the foregoing (for example, argument from physiology, bench research or first principles).

For all the outcomes included in this review, a silver level of evidence has been given, mostly because of the small samples size.

Effects of interventions

Analysis of Trials with a Separate Control Group

There was no significant heterogeneity for any comparisons, indicating that the difference between treated and control groups was consistent across trials. Each following recommendation has been graded as level silver. Since pain was measured by the visual analogue scale in three trials (Goats 1996, Hall 1994, Walker 1987), we used weighted mean differences to pool the end study results and found a statistically significant improvement in pain with a WMD -1.10, (95% CI: -1.82, -0.39). Johannsen 1994 also demonstrated an improvement in pain relief measured this time with a 0-12 scale (Johannsen 1994). This improvement was proved by a decrease of 1.00 point (95% CI: 0.23, 1.77) for the laser group relative to the placebo group. We found a clinical relevance of 28%, 25%, 19% and 22% of efficacy respectively for each of these trials. However, Goats 1996 did not find any significant difference in pain relief measured with the McGill pain questionnaire (Goats 1996). Statistically significant improvements were also found for tip to palm flexibility with a difference of 1.3 cm, (95% CI: 0.9 to 1.7 cm) (Johannsen 1994, Palmgren 1989) and morning stiffness duration with an improvement of 27 minutes, (95% CI: 3 to 52 minutes) (Goats 1996, Hall 1994, Palmgren 1989). For flexibility, we found a clinical relevance of 76% calculated with Johannsen 1994 results and 157% measured by Palmgren 1989 results. Results extracted from Goats 1996, Hall 1994 and Palmgren 1989 trials demonstrated a clinical relevance of 93%, -19% and -78%

respectively. Morning stiffness assessed in one trial had an unusually large variance (Goats 1996). Excluding this trial, the effect of laser on morning stiffness was still significant and similar in magnitude (difference of 30 minutes, 95% CI: 5.6 to 55 minutes). Also, another trial measured the morning stiffness duration by number of patients who did not improve but did not find a significant relative risk: 0.25 (95% CI: 0.03, 2.09) (Johannsen 1994) favouring the laser group relative to the placebo group.

There is conflicting evidence about range of motion. Goats 1996 found a positive effect of laser on knee range of motion after the treatment of this joint supported by a WMD of 31.80 (95% CI: 4.27, 31.80), but no significant difference at the ankle was noted with the same treatment (Goats 1996).

No other outcomes for RA were significantly affected by LLLT. These included functional status, swelling, range of motion (for metacarpal phalangeal, proximal interphalangeal and ankle joints), grip strength, and walking speed.

Only two trials (Goats 1996, Hall 1994) followed patients after the LLLT was discontinued. Both measured outcomes three months after the end of treatment. These RCTs found no significant differences between LLLT and placebo in any outcome measure at this time point.

For the subgroup analysis regarding methodologic quality, there was no difference between low (<3) and high (≥3) quality for pain reduction at the end of treatment.

For the subgroup analysis regarding joint versus nerve application, only one study applied LLLT to the nerves (Walker 1987). There was a trend towards greater pain reduction with joint/nerve application compared to joint application, but the confidence intervals for joint application (pain SMD: -0.8 (95% CI: -1.5, -0.2)) overlap those for nerve application (SMD: -1.5 (95% CI: -2.5, -0.6)). For the subgroup analysis concerning treatment duration, only one trial treated patients for more than 4 weeks (Walker 1987). The results of this analysis did not demonstrate a significant statistical difference with the longer treatment with an SMD for pain at 4 weeks of -0.4 (-0.8, +0.1) and an SMD pain at 10 weeks of -0.7 (-1.2, -0.3).

For the subgroup analysis concerning wavelength, the only outcome with data available for different wavelength comparisons was pain. There were no statistical difference between wavelengths. However, there was a trend for improved outcome with the 632 nm compared to 820 nm for pain although the confidence limits overlap [SMD 632 nm: -0.7 (95% CI: -1.2, -0.3) vs SMD 820 nm: -0.4 (95% CI: -0.8, 0.1)].

For the subgroup analysis concerning dosage, studies using a dosage of less than 3 Joules/cm² were considered low doses. This subgroup analysis was possible for VAS pain, grip strength and flexibility (tip to palm distance in cm). For pain (VAS), the low dose was significantly better than placebo (SMD: -0.8, 95% CI: -1.2, -0.4). In contrast, high dose studies found no difference between laser and placebo for pain (VAS). There was no dose effect on grip strength and flexibility (tip to palm).

Analysis of Contralateral Control Trials

The trial which used the opposite limb of the same individual as a control did not find any improvement in laser treated hands. There was no significant difference between laser and placebo for pain relief (RR 13.00, 95% CI: 0.79, 214.06). Less than half of the subjects found the lasered hand improved compared to the placebo treated hand. Relative to baseline, most patients reported similar or no changes in both laser and placebo treated hands. Specifically, the number of patients who reported this result were 11 out of 17 patients (Bliddal 1987). There was no difference in morning stiffness, mobility joint neither in laboratory tests.

DISCUSSION

In RA, this meta-analysis suggests that LLLT is effective at reducing pain relative to placebo and reducing morning stiffness. However, according to Lineker et al. 1999, duration of morning stiffness has a poor sensitivity to be considered as an outcome measure of morning stiffness (Lineker 1999). Being as morning stiffness is related to various other conditions, such as OA, it is not possible to make the difference between two conditions by comparing the duration. Definition of morning stiffness should be more specific to RA. The results show no statistical improvements for any of the following: localized swelling, muscle strength, functional status, joint tenderness or global assessments with laser treatment.

The major limitation of this meta-analysis is the heterogeneity of clinical application, including different dosages, wavelengths and types of LLLT. In addition, the results are subject to publication bias, if negative trials are not published. Furthermore, the treated joints in all but one of the RA trials were hand joints, and these results may not be generalized to other treatment sites due to different penetrations and dosages.

The possible systemic effects of laser use may be the key to understanding the improvement on pain scales. Although much of the source of inflammatory pain is peripheral (i.e., joint capsule, ligament, muscle and skin), it is within the central nervous system (CNS) that perception and interpretation of pain occurs, and particularly long-standing pain for RA (Coderre 1993). Perhaps laser irradiation positively modifies the sensory input to the CNS and provides an improvement in the perception of pain localized to the area of treatment. It has been well demonstrated that a painful stimulus in one extremity has a contralateral effect (Coderre 1993). The improved pain scores contralaterally in the studied trial provide clinical evidence of CNS involvement. Although there is not a sufficient number of studies to support this idea, the trend towards greater pain reduction when nerves as well as joints were irradiated in the study by Walker 1987, suggests that this treatment strategy should be explored to support CNS input theory.

The results suggest that improvement in overall pain relative to controls followed LLLT applied at local sites including the

metacarpophalangeal (MCP), proximal interphalangeal (PIP) and knee joints. Other work on the histologic evaluation of rheumatoid synovial membrane has shown that inflammatory changes present in the damaged synovial membrane can be reversed. Exposure to LLLT has been shown to result in: 1) anti-inflammatory and analgesic effects (Barabas 1988, Matulis 1983, Nishida 1990); 2) increased protein synthesis of rheumatoid synovial cells (Goldman 1980), 3) normalization of the permeability of the synovial membrane (Tsurko 1983) and 4) enhancement of regional microcirculation (Matulis 1983). According to Goldman 1980, the laser energy irradiates inflamed synovial cells and modifies the level of antigen-antibody complexes and helps stabilize the membranes in RA subjects (Goldman 1980). These processes support the hypothesis that LLLT has systemic effects (Prwemstay 1987, Young 1988). Several investigators have recently called for further investigation to establish that key histologic changes can be captured with our clinical measurements (Anderson 1993).

The results in this meta-analysis suggest that LLLT does not have long-lasting effects. In this review, only one trial (Walker 1987) had a treatment duration greater than four weeks. Support for a time-related effect of LLLT comes from animal model studies. Schultz and collaborators noted, in a pig in vitro cartilage model, that only after the fourth week of LLLT irradiation, a joint regeneration process began which they hypothesized to be a reactivation of the mitotic control mechanism (Schultz 1985). Furthermore, the healing of the joint surface lesion was not complete until the end of 6 weeks of application.

The subgroup analyses for dosage, wavelength and site of application indicated no significant effects of these factors on the effectiveness of LLLT in RA. However, these subgroup analyses indicated trends towards improved outcomes with longer dose duration, nerve application and shorter wavelengths. Even though there is more evidence that wavelengths not only determine penetration depth, but also specific tissue interactions (Bolagni 1985, Karu 1987). Our results do not appear to support this hypothesis. According to Seichert 1991, the wavelength effect of the two most commonly used lasers, HeNe and GaAlAs lasers, is negligible. Furthermore, "the laser light loses its coherency completely after only a few tenths of millimeters in depth" (Seichert 1991).

A subgroup analysis of the dosages in these RCTs did not reveal a relationship between the dosage and the outcomes of interest. The dosages were calculated by hand since they were not consistently reported in the articles. As previously stated, this situation shows that uniformity in reporting the dosages used is imperative (Morin 1996). The large variability in methodological quality reflects the need for standardization of conducting and reporting clinical trials in physiotherapy. In this meta-analysis, we were only able to pool certain outcomes, such as pain, morning stiffness and range of motion. The use of validated clinical measurements sensitive enough to detect changes is also needed (Anderson 1993). There is an urgent need for further well designed RCTs evaluating differ-

ent durations of treatment, nerve versus joint application, different wavelengths and different dosages. Further clinical hypothesis could be investigated for each of these aspects.

The development and use of a theoretical framework to rationalize LLLT and also to report LLLT application in clinical settings and in scientific reports is imperative (Calderhead 1991, Rothstein 1991). There is also a need for evaluation of standardized OMERACT outcomes, including joint count and physician and patient global assessment with laser treatment for RA. These outcomes are recognized as responsive to changes in RA status.

Several authors recommend further studies to elucidate the effectiveness of LLLT for RA (Beckerman 1992, King 1990, Quirion-DeGirardi 94). These studies are needed to contribute to evidence-based practice.

In summary, LLLT for RA is beneficial as a minimum of a four-week treatment with reductions in pain and morning stiffness. On the one hand, this meta-analysis found that pooled data gave some evidence of a clinical effect, but the outcomes were in conflict, and it must therefore be concluded that firm documentation of the application of LLLT in RA is not possible. Conversely, a possible clinical benefit in certain subgroups cannot be ruled out from the present meta-analysis and further large-scale studies are recommended with special attention to the findings in this meta-analysis (e.g. low versus high dose, wavelength, nerve versus joint application and treatment duration).

AUTHORS' CONCLUSIONS

Implications for practice

According to this meta-analysis, there is "silver" level of evidence that low level laser therapy decreases pain and morning stiffness in patients with rheumatoid arthritis if applied for at least 4 weeks. No side effects were reported. However, we conclude that there is insufficient data to draw firm conclusions.

Implications for research

There is a need for further RCTs evaluating different lengths of treatment, nerve vs joint application, different wavelengths and different dosages.

There is a need for evaluation of other OMERACT outcomes including joint count and physician and patient global assessment with laser treatment for RA. These outcomes are recognized as responsive to changes in RA status.

ACKNOWLEDGEMENTS

The authors are indebted to Mr. Chris Sulway and Ms. Nathalie Jean for their technical support and assistance in editing this

manuscript. Thanks to Jessie McGowan for her consultation on the search strategy.

REFERENCES

References to studies included in this review

Bliddal 1987 {published data only}

Bliddal H, Hellesen C, Ditlevsen P, Asselberghs J, Lyager L. Laser therapy of rheumatoid arthritis. *Scandinavian Journal of Rheumatology* 1987;**16**:225–228.

Goats 1996 {published data only}

Goats GC, Flett E, Hunter JA, Stirling A. Low intensity laser and phototherapy for rheumatoid arthritis. *Physiotherapy* 1996;**82**(5): 311–20.

Hall 1994 {published data only}

Hall J, Clarke AK, Elvins DM, Ring EFJ. Low level laser therapy is ineffective in the management of rheumatoid arthritis finger joints. *British Journal of Rheumatology* 1994;**33**:142–7.

Johannsen 1994 {published data only}

Johannsen B, Hauschild B, Remvig L, Johnsen V, Petersen M, Bieler T. Low energy laser therapy in rheumatoid arthritis. *Scandinavian Journal of Rheumatology* 1994;**23**:145–7.

Palmgren 1989 {published data only}

Palmgren N, Jensen GF, Kamma K, Windelin M, Colov HC. Low-power laser therapy in rheumatoid arthritis. *Lasers in Medical Science* 1989;**4**:193–6.

Walker 1987 {published data only}

Walker JB, Akhanjee LK, Cooney MM, Goldstein J, Tamayoshi S, Segal-Gidan F. Laser therapy for pain of rheumatoid arthritis. *Clinical Journal of Pain* 1987;**3**:54–9.

References to studies excluded from this review

Agambar 1992 {published data only}

Agambar L, Herbert KE, Scott DL. Low powered laser therapy for rheumatoid arthritis. *British Journal of Rheumatology* 1992;**31**(suppl 2):81.

Ammer K 2003 {published data only}

Ammer K. Conservative, Non Drug Treatment in Rheumatoid Arthritis. *Physical Medicine and Rehabilitation Kuror* 2003;**13**:13–20.

Asada K 1989 {published data only}

Asada K, Yutani Y, Shimazu A. Diode Laser Therapy for Rheumatoid Arthritis : A Clinical Evaluation of 102 Joints Treated With Low Rasctive-Level Laser Therapy (LLLT). *Laser Therapy* 1989;**1**:147–51.

Barabas 1989 {published data only}

Barabas K, Balint G, Gaspardy G, Meretey K, Pirooska E, Molnar E. Controlled clinical and experimental studies with Nd-phosphate-glass laser in patients with rheumatoid arthritis and its effect on the synovial membrane. [German] [KONTROLLIERTE KLINISCHE UND EXPERIMENTELLE UNTERSUCHUNGEN MIT ND-PHOSPHAT-GLAS-LASER BEI PATIENTEN MIT RHEUMATOID ARTHRITIS BZW. IHRE WIRKUNG AUF DIE SYN-

OVIALMEMBRANE.]. *Zeitschrift fur Physiotherapie* 1989;**41**(5): 293–6.

Day R 1999 {published data only}

Day R. Laser Modality Refresher. *SportEx Medicine* 1999;**6**:34–4.

De Bie 1998 {published data only}

De Bie, Verhagen AP, Lenssen AF, de Vet HCW, van den Wildenberg FAJM, Kootstra G, Knipschild PG. Efficacy of 904 nm laser therapy in the management of musculoskeletal disorders: a systematic review. *Physical Therapy Reviews* 1998;**3**:59–72.

Fulga C 1998 {published data only}

Fulga C. Anti-inflammatory Effect of Laser Therapy in Rheumatoid Arthritis. *Romanian Journal of Internal Medicine* 1998;**36**(3-4):273–9.

Goldman 1980 {published data only}

Goldman JA, Chiapella J, Casey H, Bass N, Graham J, McClatchey W, et al. Laser therapy of rheumatoid arthritis. *Lasers in Surgery and Medicine* 1980;**1**:93–101.

Heussler 1993 {published data only}

Heussler JK, Hinchey G, Margiotta E, Quinn R, Butler P, Martin J, et al. A double blind randomised trial of low power laser treatment in rheumatoid arthritis. *Annals of Rheumatic Diseases* 1993;**52**:703–706.

Matulis 1983 {published data only}

Matulis AA, Vasilenkaitis VV, Raistensky IL, Cheremnykh-Alexeenko EN, Gaigalene BA. Laser therapy and laseropuncture in rheumatoid arthritis, osteoarthritis deformans and psoriatic arthropathy. *Ther Arkh* 1983;**55**(7):92–7.

Miyagi 1989 {published data only}

Miyagi K. Double blind comparative study on the effect of low energy laser irradiation to rheumatoid arthritis [Japanese]. *Journal of Japanese Association of Physical Medicine Balneology & Climatology* 1989;**52**(3):117–26.

Oyamada 1988 {published data only}

Oyamada Y, Satodate R, Nishida J, Izu S, Aoki Y. A double blind study of low power He-Ne laser therapy in rheumatoid arthritis. *Optoelectronics in Medicine* 1988;**87**:747–50.

Sidorov VD 1999 {published data only}

Sidorov VD, et al. The interauricular laser therapy of rheumatoid arthritis. *Voprosy kurortologii, fizioterapii i lechebnoi fiz* 1999;**3**:35–45.

Sidorov VD 2000 {published data only}

Sidorov VD, et al. The Combined Laser Therapy of Rheumatoid Arthritis. *Voprosy Kurortologii, fizioterapii, i lechebnoi Fizicheskoi kult* 2000;**2**:13–18.

Taghawinejad 1985 {published data only}

Taghawinejad M, Frcke R. [Lasertherapie in der behandlung Kleiner Gelenke bie chronischer Polyarthritisu]. *Zeitschrift fur Physikalische Medizin* 1985;**14**:402–8.

Tsurko 1983 *{published data only}*

Tsurko VV, Muldiyarov PY, Sigidin YA. Laser therapy of rheumatoid arthritis (clinical and morphological study). *Ther. Ther Arkh* 1983; **55**:97–102.

Walker 1983 *{published data only}*

Walker J, Akhanjee LK, Cooney M. Laser therapy for pain of rheumatoid arthritis. American Society for Laser Medicine and Surgery Abstracts. American Society for Laser Medicine and Surgery, 1983.

Additional references

Anderson 1993

Anderson JJ, Chernoff MC. OMERACT: Conference on Outcome Measures in Rheumatoid Arthritis Clinical Trials. *Journal of Rheumatology* 1993;**20**:535–7.

Badley 1994

Badley EM, Rasooly I, Webster GK. Relative importance of musculoskeletal disorders as a cause of chronic health problems, disability and health care utilization: findings from the 1990 Ontario Health Survey. *Journal of Rheumatology* 1994;**21**:505–14.

Barabas 1988

Barabas K, Bakos J, Szabo DL, Sinay H, Gaspardy G, Bely M, et al. In vitro effect of neodymium phosphate glass laser irradiation on the synovial membrane (abstract). American Society for Laser Medicine and Surgery Abstracts. 1988; Vol. 8:176.

Bassler 1985

Bassler C, Datchy M, Reginster JY, et al. Human articular chondrocytes cultivated in three dimensions: effects of I.R. laser irradiation. Proceedings International Congress on Lasers in Medicine and Surgery. Bologna, 1985:381–5.

Beckerman 1992

Beckerman H, de Bie RA, Bouter LM, de Cuyper HJ, Oostendorp RAB. The efficacy of laser therapy for musculoskeletal and skin disorders: A criteria-based meta-analysis of randomized clinical trials.. *Physical Therapy* 1992;**72**:483–491.

Bolagni 1985

Bolagni L, Daviolo E, Volpi N. Effects of GaAs pulsed laser on ATP concentration and ATPase activity in vitro and in vivo. In Galletti G (Ed) Laser. Proceedings of the International Congress on Laser in Medicine and Surgery. Proceedings of the International Congress on Laser in Medicine and Surgery. Bologna: Monduzzi Editore, 1985: 47–51.

Brosseau 1994

Brosseau L, Mercille S, Quirion-de Girardi C. Le laser de classes I, II et III dans le traitement de la polyarthrite rhumatoïde et de l'arthrose. *Journal de Readaptation Médicale* 1994;**14**:107–116.

Brosseau 1998

Brosseau L, Morin M, Quirion-DeGardi C. A theoretical framework on low level laser therapy (classes I, II and III) application for the treatment of OA and RA. Proceedings of the Canadian Physiotherapy Association National Congress. Victoria: Canadian Physiotherapy Association, 1996:1.

Calderhead 1991

Calderhead RG. Watts a joule: on the importance of accurate and correct reporting of laser parameters in low reactive level laser therapy and photoactivation research. *Laser Therapy* 1991;**3**:177–82.

Ceccherelli 1989

Ceccherelli F, Altafini, Castro L, Avila GL, Ambrosio F, Giron GP. Diode laser in cervical myofascial pain: a double-blind study versus placebo. *Clinical Journal of Pain* 1989;**5**(4):301–4.

Coderre 1993

Coderre TJ, Katz J, Vaccarino AL, Melzack R. Contribution of central neuroplasticity to pathological pain: review of clinical and experimental evidence. *Pain* 1993;**52**:259–85.

Detsky 1992

Detsky AS, Naylor CD, O'Rourke K, McGeer AJ, L'Abbe KA. Incorporating variations in the quality of randomized trials into meta-analysis. *Journal of Clinical Epidemiology* 1992;**45**:255–65.

Gam 1993

Gam AN, Thorsen H, Lonnberg F. The effect of low-level laser therapy on musculoskeletal pain: a meta-analysis. *Pain* 1993;**52**:63–6.

Greenland 1985

Greenland S, Robins J. Estimation of a common effect parameter from sparse follow-up data. *Biometrics* 1985;**41**:55–68.

Helewa 1994

Helewa A, Smythe HA. Physical therapy in rheumatoid arthritis. In: Wolfe F, Pincus T editor(s). *Rheumatoid Arthritis. Pathogenesis, Assessment, Outcome and Treatment*. New York: Marcel Dekker, Inc., 1994:415–33.

Hennekens 1987

Hennekens CH, Buring JE. Measures of disease frequency and association. In: Mayrent SL editor(s). *Epidemiology in Medicine*. Boston: Boston, Little, Brown and Co., 1987.

Jadad 1996

Jadad A, Moore A, Carrol D, et al. Assessing the quality of reports of randomized trials: is blinding necessary?. *Control Clinical Trial* 1996;**17**:1–12.

Karu 1987

Karu TI. Photobiological fundamentals of low-power laser therapy. *IEEE J Quantum Electronics* 1987;**23**1:1703–17.

King 1990

King CE, Clelland JA, Knowles CJ, Jackson. Effect of helium-neon laser auriculotherapy on experimental pain threshold. *Physical Therapy* 1990;**70**:24–30.

Kudoh 1989

Kudoh C, Inomata K, Okajima K, Motegi M, Ohshiro T. Effects of 830nm gallium aluminium arsenide diode laser radiation on rat saphenous nerve sodium-potassium-adenosine triphosphate activity: A possible pain attenuation mechanism explained. *Laser Therapy* 1989;**1**:63–7.

Lineker 1999

Lineker S, Badley E, Charles C, Hart L, Streiner D. Defining Morning Stiffness in Rheumatoid Arthritis. *Journal of Rheumatology* 1999; **29**:1052–7.

Mantel 1959

Mantel N, Haenszel W. Statistical aspects of the analysis of data from retrospective studies of disease. *Journal of the National Cancer Institute* 1959;**22**:719–48.

Morin 1996

Morin M, Brosseau L, Quirion-DeGardi C. A theoretical framework on low level laser therapy (classes I, II and III) application for the

treatment of OA and RA. Proceedings of the Canadian Physiotherapy Association National Congress, Victoria (B.C. Proceedings of the Canadian Physiotherapy Association National Congress. Canadian Physiotherapy Association, 1996:1.

Nishida 1990

Nishida J, Satoh T, Satodate R, Abe M, Oyamada Y. Histological evaluation of the effect of HeNe laser irradiation on the synovial membrane in rheumatoid arthritis. *Japanese Journal of Rheumatology* 1990;2:251–60.

OMERACT 1993

OMERACT Proceedings. Conference on Outcome Measures in Rheumatoid Arthritis Clinical Trials. *Journal of Rheumatology* 1993; 20:526–91.

Petitti 1994

Petitti D. Meta-analysis, decision analysis, and cost-effectiveness analysis: methods for quantitative synthesis in medicine. *Statistics in Medicine*. New York: Oxford University Press, 1994:90–114.

Prwemstay 1987

Prwemstay C, Davis CC. Interaction of electromagnetic fields with genetic information. In: M Balnk, E Finkl editor(s). *Mechanistic approaches to interactions of electric and electromagnetic fields with living system*. New York: Planum Press, 1987:231.

Quirion-DeGirardi 94

Quirion-DeGirardi C, Brosseau L, Mercille S. Le laser de classes I, II et III dans le traitement de la polyarthrite rhumatoïde et de l'arthrose:

Partie 2 Recension des études cliniques. *JRéadapt Méd* 1994;14: 139–46.

Rothstein 1991

Rothstein JM. Theoretically speaking. *Physical Therapy* 1991;11: 789–90.

Schultz 1985

Schultz RJ, Krishnamurthy S, Thelmo W, Rodriguez JE, Harvey G. Effects of varying intensities of laser energy on articular cartilage: A preliminary study. *Lasers in Surgery and Medicine* 1985;5:577–88.

Seichert 1991

Seichert N. Controlled trials of laser treatment. In (eds): Physiotherapy: Controlled Trials and Facts. Rheumatology [Physiotherapy: Controlled Trials and Facts. Rheumatology]. In: Schlapbach P, Gerber NJ editor(s). *Basel, Karger*,. Vol. 14, Basel: Karger, 1991:205–17.

Tugwell 2004

Tugwell P, Shea B, Boers M, Brooks P, Strand V, et al. *Evidence Based Rheumatology*. BMJ Books, 2004.

Walker 1996

Walker JM, Helewa A. *Physical Therapy in Arthritis*. Philadelphia: W.B. Saunders Co, 1996.

Young 1988

Young S, Harvey W, Dyson M, Diamantopoulos C. Macrophage responsiveness to laser therapy. *Lasers in Surgery and Medicine* 1988; 8:186–90.

* Indicates the major publication for the study

CHARACTERISTICS OF STUDIES

Characteristics of included studies [ordered by study ID]

Bliddal 1987

Methods	Randomized Double-blind Placebo Sample size at entry 17 Patients being their own control Study duration: 3 weeks and follow-up at 4 weeks
Participants	Patients with active RA and symmetrical involvement of the metacarpophalangeal joint of the index. No change in other therapy was accepted during the study. F/M: 16/2 Median age: 57 years Median duration of RA: 10 years Class II: 11 Class III: 7

Bliddal 1987 (Continued)

Interventions	<p>Patients were randomized to therapy on the MP joint of the right and left index with laser or placebo. Therapy was given on 3 alternate days a week for 3 consecutive weeks followed by an observation time of 4 weeks. Laser: He-Ne Wavelength: 633nm Output power: 10mW Spot size : 0.5cm² Dosage:6J/cm² Irradiation time: 5 min Placebo: Placebo apparatus containing a red 12 V, 10 W bulb and a reflecting mirror giving as far as possible the same red light through an identical optic fibre and pen</p>
Outcomes	<p>Pain (VAS) Morning stiffness</p>
Notes	<p>Quality Jadad's 4/5</p>

Goats 1996

Methods	<p>Randomized Double blind (partial) Placebo controlled Sample size at entry: LAser: 25; placebo: 10 Study duration: 6 months</p>
Participants	<p>Patients with RA affecting 2 or more of tibio-femoral, talocrural, subtalar, midtarsal or MCP joints No steroids, immunosuppressants or intra-articular injections in the last 2 months Mean age (yrs): Laser 57 yrs, placebo 64 yrs Sex % F: Laser 80%; placebo: 80% Disease duration (mean): Laser 7.54 yrs, Placebo: 9.80 yrs</p>
Interventions	<p>Gr1: Laser: Ga-As-Al Class: 3B Wavelength: 850 nm Output power: 940 mW Power density: Frequency: 5 Hz Duty cycle: 80% Spot size : 0.125 cm² Divergence: Dosage: 8.1 J/cm² to each aspect of the joint (24.3 J/cm² for each right and left tibiofemoral joint, 16.2 J/cm² for each right and left MTP) Area treated: Tibiofemoral, talocrural, subtalar, midtarsal and metatarsophalangeal joints Irradiation time: 4 minutes per joint Schedule: 2 times/week for 4 weeks Total sessions: 8 Calibrated: no</p>

Goats 1996 (Continued)

	Placebo: Dummy laser probe, subjects wore eye protection that dimmed visible radiation Measurements: 0, 1, 3, and 6 months after start of treatment Gr2:Gr 2 : Placebo (identical in external appearance but having no output).
Outcomes	Pain (VAS 10 cm) McGill Pain questionnaire ESR CRP HAQ Hemoglobin Platelets Knee ROM Ankle ROM Suprapatellar swelling (cm) Morning stiffness duration (hours) Walking speed (sec) Ritchie index Rheumatoid factor
Notes	Quality Jadad's: 2/5

Hall 1994

Methods	Randomized Triple blind Sample size at entry: Laser 20; placebo : 20 Study duration: 6 months
Participants	Patients with RA, Steinbrocker functional class II or III, and active synovitis of some or all of the MCP and PIP joints No changes to NSAIDs in last 30 days, DMARDS last 3 months, joints incapable of response for mechanical reasons Setting: outpatient clinics Mean age (yrs): Laser: 67.1; Placebo: 60.9 Sex % F: Laser 85%; Placebo 85% Disease duration: Laser: 146.4 months, Placebo: 111.6 months Duration of hand symptoms: Laser 52.5 months; placebo: 54.5 months
Interventions	Laser: Ga-As-Al Wavelength: 820 nm Output power: Actual 40 mW (Rated 50 mW) Power density: Frequency: 5 Hz Duty cycle: 80% Spot size : 0.1 cm ²

Hall 1994 (Continued)

	Divergence: 6 degrees Dosage: 3.6 J/cm ² per radiant exposure for each joints Area treated: 90 seconds per joint on the radial, ulnar, dorsal, ventral aspects of 1st to 5th MCP and PIP joints of the most affected hand Irradiation time: 18 minutes per treatment Schedule: 4 weeks Total sessions: 12 Calibrated: no Placebo: Dummy laser probe (identical apparatus)
Outcomes	Pain (10 cm VAS) Health Assessment Questionnaire (HAQ) Grip strength (mmHg) Hemoglobin (g/dL) CRP (g/mL) Ritchie Index Platelets (x10/L) Duration of morning stiffness (minutes) MCP swelling (cm) PIP swelling (cm) MCP ROM (degrees) PIP ROM (degrees) Thermographic index
Notes	Quality Jadad's: 3/5

Johannsen 1994

Methods	Randomized Double-blind Placebo controlled Sample size at entry: Laser: 10 Placebo: 12 Study duration 4 weeks
Participants	Participants with RA, Steinbrocker functional class I or II With no changes to steroids, NSAID or analgesics in last month and no changes to DMARDs within the last 3 months Median age : Laser 59 yrs, Placebo 62 yrs Sex % F: Laser: 90; placebo: 83 Disease duration not measured

Johannsen 1994 (Continued)

Interventions	<p>Laser: GaAsAl Wavelength: 830 nm Output power: 21 mW Power density: NA Frequency: NA Duty cycle: NA Spot size: 0.07 cm² spot size Divergence: NA Dosage: 23.2 J per treatment with 2.9J on 4 points on 2 joints Area treated: 2 most painful metacarpal phalangeal joints (MCP) on worst affected hand Schedule: 3 times/week for 4 weeks Total sessions: 12 Calibration: yes Placebo: Same apparatus giving red visible light</p>
Outcomes	<p>Joint tenderness Pain (range 0 to 12) Erythrocyte sedimentation rate (ESR) C Reactive Protein (CRP) Pinch strength Flexibility- tip to palm distance (cm) Grip strength (kg) Improvement in morning stiffness</p>
Notes	<p>Quality Jadad's: 5/5</p>

Palmgren 1989

Methods	<p>Randomized Double-blind Placebo controlled Sample size at entry: Laser : 19; placebo : 16 Study duration: 4 weeks</p>
Participants	<p>Patients with classic RA, Steinbrocker functional class between I and III No changes to basic medication during study Patients with unstable disease excluded Mean age (yrs) : Laser, M: 66.0, Laser, F: 61.1; Placebo M: 68.0, Placebo F: 57.5 Sex % F: Laser 74%; Placebo: 88% Disease duration: Laser: 13.4 yrs; Placebo: 15.5 yrs</p>
Interventions	<p>Laser: GaAlAs Wavelength: 820 nm Output power: 15 mW Power density: NA</p>

Palmgren 1989 (Continued)

	<p>Frequency: NA Duty cycle: NA Spot size: 0.1256 cm² Divergence: 6 degrees Dosage: 5.5 J/cm² Irradiation time: 60 s on each lateral side of each joint Area treated: 2nd to 5th MCP and proximal interphalangeal (PIP) joints of the most affected hand Schedule: 3 times/week for 4 weeks Total sessions: 12 Placebo: Similar but disconnected diode</p>
Outcomes	<p>Joint swelling ESR Fibrinogen (mmol/L) Leukocytes (10⁹/L) Lymphocytes (%) Hemoglobin (mmol/L) Grip strength Flexibility - tip to palm distance (mm) Duration morning stiffness (hrs)</p>
Notes	Quality Jadad's: 3/5

Walker 1987

Methods	<p>Partially blinded Placebo controlled Sample size at entry: Laser : 38; Placebo : 34 Study duration 10 weeks</p>
Participants	<p>Participants with RA, defined by the American Rheumatism Association (ARA) Median age: Laser 61.5 yrs, Placebo 60.0 yrs Sex % F: Laser 60%, Placebo 79% Disease duration (median yrs): Laser 6 yrs, Placebo 11 yrs</p>
Interventions	<p>Laser: He-Ne Class: I Wavelength: 632.5 nm Output power: 1 mW Power density: 47.7 per cm² Frequency: 20 Hz Duty cycle: 50% Spot size: 4 mm² Divergence: 30 degrees Dosage: 3.6 mJ/cm²</p>

Walker 1987 (Continued)

	Area treated: Radial, median, saphenous nerves for 20 s bilaterally and painful joints for 3 x 10 s along the circumference Schedule: 3 times/week for 10 weeks Total sessions: 30 Placebo: Sham apparatus with no light, subjects closed eyes
Outcomes	Pain (10 cm visual analogue scale)
Notes	Quality Jadad's: 1/5

Characteristics of excluded studies [ordered by study ID]

Agambar 1992	Only abstract available
Ammer K 2003	Literature Review
Asada K 1989	No control group
Barabas 1989	Foreign language
Day R 1999	Literature Review
De Bie 1998	Literature Review
Fulga C 1998	Patients with juvenile RA; no control group
Goldman 1980	Insufficient statistical data
Heussler 1993	Insufficient statistical data
Matulis 1983	No control group
Miyagi 1989	Foreign language
Oyamada 1988	Although reported as double-blind in the abstract, a reply from the authors indicated that this trial was not randomized
Sidorov VD 1999	Abstract only available in English; Article was in Russian
Sidorov VD 2000	Abstract only available in English; Article was in Russian

(Continued)

Taghawinejad 1985	Foreign language
Tsurko 1983	Insufficient statistical data
Walker 1983	Duplicate of Walker 1987

DATA AND ANALYSES

Comparison 1. Laser vs Placebo- End of treatment (approx 10 wks)

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Pain (Visual Analogue Scale 10 cm)	3	147	Mean Difference (IV, Fixed, 95% CI)	-1.10 [-1.82, -0.39]
2 Pain (0-12 scale)	1	22	Mean Difference (IV, Fixed, 95% CI)	-1.01 [-1.77, -0.23]
3 McGill Pain questionnaire	1	35	Mean Difference (IV, Fixed, 95% CI)	1.06 [-1.22, 4.22]
4 Ritchie Index	1	40	Mean Difference (IV, Fixed, 95% CI)	2.60 [-4.50, 9.70]
5 Health Assessment Questionnaire (HAQ)	2	75	Mean Difference (IV, Fixed, 95% CI)	0.93 [-2.01, 3.87]
6 MCP ROM (degrees)	1	40	Mean Difference (IV, Fixed, 95% CI)	-0.06 [-15.99, 14.99]
7 PIP ROM (degrees)	1	40	Mean Difference (IV, Fixed, 95% CI)	4.01 [-6.60, 14.60]
8 Knee ROM	1	70	Mean Difference (IV, Fixed, 95% CI)	-18.03 [-31.80, -4.27]
8.1 Right knee ROM	1	35	Mean Difference (IV, Fixed, 95% CI)	-12.90 [-31.98, 6.18]
8.2 Left knee ROM	1	35	Mean Difference (IV, Fixed, 95% CI)	-23.60 [-43.47, -3.73]
9 Ankle ROM	1	70	Mean Difference (IV, Fixed, 95% CI)	4.60 [-0.33, 9.53]
9.1 Right ankle ROM	1	35	Mean Difference (IV, Fixed, 95% CI)	5.90 [-1.21, 13.01]
9.2 Left ankle ROM	1	35	Mean Difference (IV, Fixed, 95% CI)	3.40 [-3.43, 10.23]
10 Flexibility- tip to palm distance (cm)	2	57	Mean Difference (IV, Fixed, 95% CI)	-1.28 [-1.72, -0.85]
11 Morning stiffness duration (min)	3	110	Mean Difference (IV, Fixed, 95% CI)	-27.45 [-51.95, -2.95]
12 Morning stiffness not improved	1	22	Risk Ratio (M-H, Fixed, 95% CI)	0.76 [0.49, 1.19]
13 Rheumatoid factor positive	1	35	Risk Ratio (M-H, Fixed, 95% CI)	0.10 [0.61, 1.34]
14 Grip strength	3	97	Mean Difference (IV, Fixed, 95% CI)	0.62 [-0.36, 1.60]
14.1 Grip strength (mmHg)	2	75	Mean Difference (IV, Fixed, 95% CI)	7.71 [0.15, 15.27]
14.2 Grip strength (kg)	1	22	Mean Difference (IV, Fixed, 95% CI)	0.06 [-0.49, 1.49]
15 Suprapatellar swelling - right knee (cm)	1	35	Mean Difference (IV, Fixed, 95% CI)	-2.94 [-6.31, 0.43]
16 Suprapatellar swelling- left knee (cm)	1	35	Mean Difference (IV, Fixed, 95% CI)	-2.69 [-6.33, 0.95]
17 MCP swelling (cm)	1	40	Mean Difference (IV, Fixed, 95% CI)	0.20 [-0.78, 1.18]
18 PIP swelling (cm)	2	75	Mean Difference (IV, Fixed, 95% CI)	0.53 [-1.12, 2.18]
19 Walking speed (sec)	1	35	Mean Difference (IV, Fixed, 95% CI)	-1.87 [-7.08, 3.34]
20 Fibrinogen	1	35	Mean Difference (IV, Fixed, 95% CI)	1.06 [-0.00, 3.00]
21 Leukocytes	1	35	Mean Difference (IV, Fixed, 95% CI)	1.60 [0.62, 2.58]
22 Lymphocytes	1	35	Mean Difference (IV, Fixed, 95% CI)	3.01 [-6.57, 12.57]
23 ESR (mm/hr)	3	92	Mean Difference (IV, Fixed, 95% CI)	-10.09 [-15.04, -5.15]
24 CRP (g/mL)	2	57	Mean Difference (IV, Fixed, 95% CI)	2.55 [-14.11, 19.21]
25 Hemoglobin	2	70	Mean Difference (IV, Fixed, 95% CI)	0.47 [0.01, 0.93]

26 Platelets	1	35	Mean Difference (IV, Fixed, 95% CI)	20.35 [-64.36, 105.06]
--------------	---	----	-------------------------------------	------------------------

Comparison 2. Laser vs Placebo- End of follow-up (approx 20 wks)

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Pain	2	54	Mean Difference (IV, Fixed, 95% CI)	0.06 [-1.24, 1.36]
1.1 Pain (10 cm visual analogue scale)	2	54	Mean Difference (IV, Fixed, 95% CI)	0.06 [-1.24, 1.36]
2 McGill Pain questionnaire	1	28	Mean Difference (IV, Fixed, 95% CI)	1.39 [-2.23, 5.01]
3 Health Assessment Questionnaire (HAQ)	2	54	Mean Difference (IV, Fixed, 95% CI)	1.10 [-2.80, 4.99]
4 Ritchie Index	1	26	Mean Difference (IV, Fixed, 95% CI)	3.30 [-5.74, 12.34]
5 PIP ROM (degrees)	1	26	Mean Difference (IV, Fixed, 95% CI)	2.01 [-16.30, 20.30]
6 Knee ROM	1	56	Mean Difference (IV, Fixed, 95% CI)	-11.47 [-31.26, 8.31]
6.1 Right knee ROM	1	28	Mean Difference (IV, Fixed, 95% CI)	-13.60 [-41.44, 14.24]
6.2 Left knee ROM	1	28	Mean Difference (IV, Fixed, 95% CI)	-9.30 [-37.43, 18.83]
7 Ankle ROM	1	56	Mean Difference (IV, Fixed, 95% CI)	-1.20 [-11.69, 9.28]
7.1 Right ankle ROM	1	28	Mean Difference (IV, Fixed, 95% CI)	1.80 [-12.59, 16.19]
7.2 Left ankle ROM	1	28	Mean Difference (IV, Fixed, 95% CI)	-4.60 [-19.91, 10.71]
8 MCP ROM (degrees)	1	26	Mean Difference (IV, Fixed, 95% CI)	10.01 [-6.86, 26.86]
9 Morning stiffness duration (min)	2	54	Mean Difference (IV, Fixed, 95% CI)	12.16 [-31.00, 55.31]
10 Walking speed (sec)	1	28	Mean Difference (IV, Fixed, 95% CI)	-1.91 [-12.60, 8.78]
11 Grip strength	1	26	Mean Difference (IV, Fixed, 95% CI)	-29.01 [-61.44, 3.44]
11.1 Grip strength (mmHg)	1	26	Mean Difference (IV, Fixed, 95% CI)	-29.01 [-61.44, 3.44]
11.2 Grip strength (kg)	0	0	Mean Difference (IV, Fixed, 95% CI)	Not estimable
12 Suprapatellar swelling- left knee (cm)	1	28	Mean Difference (IV, Fixed, 95% CI)	-2.89 [-14.38, 8.60]
13 Suprapatellar swelling - right knee (cm)	1	28	Mean Difference (IV, Fixed, 95% CI)	-3.29 [-14.66, 8.08]
14 MCP swelling (cm)	1	26	Mean Difference (IV, Fixed, 95% CI)	-0.40 [-1.23, 0.43]
15 PIP swelling (cm)	1	26	Mean Difference (IV, Fixed, 95% CI)	0.90 [-1.88, 3.68]
16 Thermographic index	1	26	Mean Difference (IV, Fixed, 95% CI)	0.60 [-0.11, 1.31]
17 Rheumatoid factor positive	1	20	Risk Ratio (M-H, Fixed, 95% CI)	1.01 [0.60, 1.68]
18 ESR (mm/hr)	1	28	Mean Difference (IV, Fixed, 95% CI)	-2.01 [-26.94, 22.94]
19 CRP (g/mL)	2	54	Mean Difference (IV, Fixed, 95% CI)	-4.21 [-20.25, 11.83]
20 Hemoglobin	2	54	Mean Difference (IV, Fixed, 95% CI)	0.98 [-0.21, 2.16]

21 Platelets	2	54	Mean Difference (IV, Fixed, 95% CI)	32.68 [-37.76, 103.11]
--------------	---	----	-------------------------------------	------------------------

Comparison 3. Treatment length- subgroup analysis for end of treatment results

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Pain (10 cm VAS or 0-12 scale)	4		Mean Difference (IV, Fixed, 95% CI)	Subtotals only
1.1 Treatment for 4 weeks	3	97	Mean Difference (IV, Fixed, 95% CI)	-0.83 [-1.46, -0.20]
1.2 Treatment for 10 weeks	1	72	Mean Difference (IV, Fixed, 95% CI)	-1.55 [-2.47, -0.61]

Comparison 4. Treatment length- subgroup analysis for end of follow up results

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Pain (10 cm VAS or 0-12 scale)	2		Mean Difference (IV, Fixed, 95% CI)	Subtotals only
1.1 Treatment for 4 weeks	2	54	Mean Difference (IV, Fixed, 95% CI)	0.06 [-1.24, 1.36]
1.2 Treatment for 10 weeks	0	0	Mean Difference (IV, Fixed, 95% CI)	Not estimable

Comparison 5. Methodologic quality: low (<3) vs high (=> 3)

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Pain at end of treatment	4	169	Mean Difference (IV, Fixed, 95% CI)	-1.05 [-1.58, -0.53]
1.1 Low quality (<3)	2	107	Mean Difference (IV, Fixed, 95% CI)	-1.39 [-2.23, -0.56]
1.2 High quality (=> 3)	2	62	Mean Difference (IV, Fixed, 95% CI)	-0.84 [-1.51, -0.16]

Comparison 6. Laser vs Placebo - Joint vs Nerve - end of treatment

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Pain (10 cm VAS or 0-12 scale)	4		Mean Difference (IV, Fixed, 95% CI)	Subtotals only
1.1 Joint application	3	97	Mean Difference (IV, Fixed, 95% CI)	-0.83 [-1.46, -0.20]
1.2 Nerve application	0	0	Mean Difference (IV, Fixed, 95% CI)	Not estimable
1.3 Joint and nerve application	1	72	Mean Difference (IV, Fixed, 95% CI)	-1.55 [-2.47, -0.61]

Comparison 7. Laser vs Placebo - Joint vs Nerve - end of follow up

Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis (Review)
Copyright © 2009 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Pain (10 cm VAS or 0-12 scale)	2		Mean Difference (IV, Fixed, 95% CI)	Subtotals only
1.1 Joint application	2	54	Mean Difference (IV, Fixed, 95% CI)	0.06 [-1.24, 1.36]
1.2 Nerve application	0	0	Mean Difference (IV, Fixed, 95% CI)	Not estimable
1.3 Joint and nerve application	0	0	Mean Difference (IV, Fixed, 95% CI)	Not estimable

Comparison 8. Laser vs Placebo- Wavelength analysis - End of treatment

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Pain	4		Mean Difference (IV, Fixed, 95% CI)	Subtotals only
1.1 Wavelength < 660 nm	1	72	Mean Difference (IV, Fixed, 95% CI)	-1.55 [-2.47, -0.61]
1.2 Wavelength 820, 830 nm	2	62	Mean Difference (IV, Fixed, 95% CI)	-0.84 [-1.51, -0.16]
1.3 Wavelength 650-950 nm	1	35	Mean Difference (IV, Fixed, 95% CI)	-0.78 [-2.68, 1.12]
2 McGill Pain questionnaire	1		Mean Difference (IV, Fixed, 95% CI)	Subtotals only
2.1 Wavelength < 660 nm	0	0	Mean Difference (IV, Fixed, 95% CI)	Not estimable
2.2 Wavelength 820, 830 nm	0	0	Mean Difference (IV, Fixed, 95% CI)	Not estimable
2.3 Wavelength 650-950 nm	1	35	Mean Difference (IV, Fixed, 95% CI)	1.06 [-1.22, 4.22]
3 Ritchie Index	1		Mean Difference (IV, Fixed, 95% CI)	Subtotals only
3.1 Wavelength < 660 nm	0	0	Mean Difference (IV, Fixed, 95% CI)	Not estimable
3.2 Wavelength 820, 830 nm	1	40	Mean Difference (IV, Fixed, 95% CI)	2.60 [-4.50, 9.70]
3.3 Wavelength 650-950 nm	0	0	Mean Difference (IV, Fixed, 95% CI)	Not estimable
4 Health Assessment Questionnaire (HAQ)	2		Mean Difference (IV, Fixed, 95% CI)	Subtotals only
4.1 Wavelength < 660 nm	0	0	Mean Difference (IV, Fixed, 95% CI)	Not estimable
4.2 Wavelength 820, 830 nm	1	40	Mean Difference (IV, Fixed, 95% CI)	1.26 [-2.09, 4.59]
4.3 Wavelength 650-950 nm	1	35	Mean Difference (IV, Fixed, 95% CI)	-0.18 [-6.39, 6.03]
5 MCP ROM (degrees)	1		Mean Difference (IV, Fixed, 95% CI)	Subtotals only
5.1 Wavelength < 660 nm	0	0	Mean Difference (IV, Fixed, 95% CI)	Not estimable
5.2 Wavelength 820, 830 nm	1	40	Mean Difference (IV, Fixed, 95% CI)	-0.06 [-15.99, 14.99]
5.3 Wavelength 650-950 nm	0	0	Mean Difference (IV, Fixed, 95% CI)	Not estimable
6 PIP ROM (degrees)	1		Mean Difference (IV, Fixed, 95% CI)	Subtotals only
6.1 Wavelength < 660 nm	0	0	Mean Difference (IV, Fixed, 95% CI)	Not estimable
6.2 Wavelength 820, 830 nm	1	40	Mean Difference (IV, Fixed, 95% CI)	4.01 [-6.60, 14.60]
6.3 Wavelength 650-950 nm	0	0	Mean Difference (IV, Fixed, 95% CI)	Not estimable
7 Left knee Range of motion	1		Mean Difference (IV, Fixed, 95% CI)	Subtotals only
7.1 Wavelength < 660 nm	0	0	Mean Difference (IV, Fixed, 95% CI)	Not estimable
7.2 Wavelength 820, 830 nm	0	0	Mean Difference (IV, Fixed, 95% CI)	Not estimable
7.3 Wavelength 650-950 nm	1	35	Mean Difference (IV, Fixed, 95% CI)	-23.60 [-43.47, -3.73]
8 Right knee ROM	1		Mean Difference (IV, Fixed, 95% CI)	Subtotals only
8.1 Wavelength < 660 nm	0	0	Mean Difference (IV, Fixed, 95% CI)	Not estimable
8.2 Wavelength 820, 830 nm	0	0	Mean Difference (IV, Fixed, 95% CI)	Not estimable

8.3 Wavelength 650-950 nm	1	35	Mean Difference (IV, Fixed, 95% CI)	-12.90 [-31.98, 6.18]
9 Left ankle ROM	1		Mean Difference (IV, Fixed, 95% CI)	Subtotals only
9.1 Wavelength < 660 nm	0	0	Mean Difference (IV, Fixed, 95% CI)	Not estimable
9.2 Wavelength 820, 830 nm	0	0	Mean Difference (IV, Fixed, 95% CI)	Not estimable
9.3 Wavelength 650-950 nm	1	35	Mean Difference (IV, Fixed, 95% CI)	3.40 [-3.43, 10.23]
10 Right ankle ROM	1		Mean Difference (IV, Fixed, 95% CI)	Subtotals only
10.1 Wavelength < 660 nm	0	0	Mean Difference (IV, Fixed, 95% CI)	Not estimable
10.2 Wavelength 820, 830 nm	0	0	Mean Difference (IV, Fixed, 95% CI)	Not estimable
10.3 Wavelength 650-950 nm	1	35	Mean Difference (IV, Fixed, 95% CI)	7.90 [0.79, 15.01]
11 Flexibility- tip to palm distance (cm)	2		Mean Difference (IV, Fixed, 95% CI)	Subtotals only
11.1 Wavelength < 660 nm	0	0	Mean Difference (IV, Fixed, 95% CI)	Not estimable
11.2 Wavelength 820, 830 nm	2	57	Mean Difference (IV, Fixed, 95% CI)	-1.28 [-1.72, -0.85]
11.3 Wavelength 650-950 nm	0	0	Mean Difference (IV, Fixed, 95% CI)	Not estimable
12 Morning stiffness duration (min)	3		Mean Difference (IV, Fixed, 95% CI)	Subtotals only
12.1 Wavelength < 660 nm	0	0	Mean Difference (IV, Fixed, 95% CI)	Not estimable
12.2 Wavelength 820, 830 nm	2	75	Mean Difference (IV, Fixed, 95% CI)	-30.65 [-55.71, -5.58]
12.3 Wavelength 650-950 nm	1	35	Mean Difference (IV, Fixed, 95% CI)	40.80 [-75.05, 156.65]
13 Morning stiffness not improved	1		Risk Ratio (M-H, Fixed, 95% CI)	Subtotals only
13.1 Wavelength < 660 nm	0	0	Risk Ratio (M-H, Fixed, 95% CI)	Not estimable
13.2 Wavelength 820, 830 nm	1	22	Risk Ratio (M-H, Fixed, 95% CI)	0.76 [0.49, 1.19]
13.3 Wavelength 650-950 nm	0	0	Risk Ratio (M-H, Fixed, 95% CI)	Not estimable
14 Rheumatoid factor positive	1		Risk Ratio (M-H, Fixed, 95% CI)	Subtotals only
14.1 Wavelength < 660 nm	0	0	Risk Ratio (M-H, Fixed, 95% CI)	Not estimable
14.2 Wavelength 820, 830 nm	0	0	Risk Ratio (M-H, Fixed, 95% CI)	Not estimable
14.3 Wavelength 650-950 nm	1	35	Risk Ratio (M-H, Fixed, 95% CI)	0.10 [0.61, 1.34]
15 Grip strength (mmHg, KPa, kg)	3		Mean Difference (IV, Fixed, 95% CI)	Subtotals only
15.1 Wavelength < 660 nm	0	0	Mean Difference (IV, Fixed, 95% CI)	Not estimable
15.2 Wavelength 820, 830 nm	3	97	Mean Difference (IV, Fixed, 95% CI)	0.62 [-0.36, 1.60]
15.3 Wavelength 650-950 nm	0	0	Mean Difference (IV, Fixed, 95% CI)	Not estimable
16 Suprapatellar swelling - right knee (cm)	1		Mean Difference (IV, Fixed, 95% CI)	Subtotals only
16.1 Wavelength < 660 nm	0	0	Mean Difference (IV, Fixed, 95% CI)	Not estimable
16.2 Wavelength 820, 830 nm	0	0	Mean Difference (IV, Fixed, 95% CI)	Not estimable
16.3 Wavelength 650-950 nm	1	35	Mean Difference (IV, Fixed, 95% CI)	-2.94 [-6.31, 0.43]
17 Suprapatellar swelling- left knee (cm)	1		Mean Difference (IV, Fixed, 95% CI)	Subtotals only
17.1 Wavelength < 660 nm	0	0	Mean Difference (IV, Fixed, 95% CI)	Not estimable
17.2 Wavelength 820, 830 nm	0	0	Mean Difference (IV, Fixed, 95% CI)	Not estimable
17.3 Wavelength 650-950 nm	1	35	Mean Difference (IV, Fixed, 95% CI)	-2.69 [-6.33, 0.95]
18 MCP swelling (cm)	1		Mean Difference (IV, Fixed, 95% CI)	Subtotals only
18.1 Wavelength < 660 nm	0	0	Mean Difference (IV, Fixed, 95% CI)	Not estimable
18.2 Wavelength 820, 830 nm	1	40	Mean Difference (IV, Fixed, 95% CI)	0.20 [-0.78, 1.18]
18.3 Wavelength 650-950 nm	0	0	Mean Difference (IV, Fixed, 95% CI)	Not estimable
19 PIP swelling (cm)	2		Mean Difference (IV, Fixed, 95% CI)	Subtotals only
19.1 Wavelength < 660 nm	0	0	Mean Difference (IV, Fixed, 95% CI)	Not estimable
19.2 Wavelength 820, 830 nm	2	75	Mean Difference (IV, Fixed, 95% CI)	0.53 [-1.12, 2.18]

19.3 Wavelength 650-950 nm	0	0	Mean Difference (IV, Fixed, 95% CI)	Not estimable
20 Walking speed (sec)	1		Mean Difference (IV, Fixed, 95% CI)	Subtotals only
20.1 Wavelength < 660 nm	0	0	Mean Difference (IV, Fixed, 95% CI)	Not estimable
20.2 Wavelength 820, 830 nm	0	0	Mean Difference (IV, Fixed, 95% CI)	Not estimable
20.3 Wavelength 650-950 nm	1	35	Mean Difference (IV, Fixed, 95% CI)	-1.87 [-7.08, 3.34]
21 Fibrinogen	1		Mean Difference (IV, Fixed, 95% CI)	Subtotals only
21.1 Wavelength < 660 nm	0	0	Mean Difference (IV, Fixed, 95% CI)	Not estimable
21.2 Wavelength 820, 830 nm	1	35	Mean Difference (IV, Fixed, 95% CI)	1.06 [-0.00, 3.00]
21.3 Wavelength 650-950 nm	0	0	Mean Difference (IV, Fixed, 95% CI)	Not estimable
22 Leukocytes	1		Mean Difference (IV, Fixed, 95% CI)	Subtotals only
22.1 Wavelength < 660 nm	0	0	Mean Difference (IV, Fixed, 95% CI)	Not estimable
22.2 Wavelength 820, 830 nm	1	35	Mean Difference (IV, Fixed, 95% CI)	1.60 [0.62, 2.58]
22.3 Wavelength 650-950 nm	0	0	Mean Difference (IV, Fixed, 95% CI)	Not estimable
23 Lymphocytes	1		Mean Difference (IV, Fixed, 95% CI)	Subtotals only
23.1 Wavelength < 660 nm	0	0	Mean Difference (IV, Fixed, 95% CI)	Not estimable
23.2 Wavelength 820, 830 nm	1	35	Mean Difference (IV, Fixed, 95% CI)	3.01 [-6.57, 12.57]
23.3 Wavelength 650-950 nm	0	0	Mean Difference (IV, Fixed, 95% CI)	Not estimable
24 ESR (mm/hr)	3		Mean Difference (IV, Fixed, 95% CI)	Subtotals only
24.1 Wavelength < 660 nm	0	0	Mean Difference (IV, Fixed, 95% CI)	Not estimable
24.2 Wavelength 820, 830 nm	2	57	Mean Difference (IV, Fixed, 95% CI)	-11.12 [-16.21, -6.04]
24.3 Wavelength 650-950 nm	1	35	Mean Difference (IV, Fixed, 95% CI)	8.27 [-13.18, 29.72]
25 CRP (g/mL)	2		Mean Difference (IV, Fixed, 95% CI)	Subtotals only
25.1 Wavelength < 660 nm	0	0	Mean Difference (IV, Fixed, 95% CI)	Not estimable
25.2 Wavelength 820, 830 nm	1	22	Mean Difference (IV, Fixed, 95% CI)	-120.01 [-190.13, -49.87]
25.3 Wavelength 650-950 nm	1	35	Mean Difference (IV, Fixed, 95% CI)	9.88 [-7.27, 27.03]
26 Hemoglobin	2		Mean Difference (IV, Fixed, 95% CI)	Subtotals only
26.1 Wavelength < 660 nm	0	0	Mean Difference (IV, Fixed, 95% CI)	Not estimable
26.2 Wavelength 820, 830 nm	1	35	Mean Difference (IV, Fixed, 95% CI)	0.60 [0.11, 1.09]
26.3 Wavelength 650-950 nm	1	35	Mean Difference (IV, Fixed, 95% CI)	-0.69 [-2.12, 0.74]
27 Platelets	1		Mean Difference (IV, Fixed, 95% CI)	Subtotals only
27.1 Wavelength < 660 nm	0	0	Mean Difference (IV, Fixed, 95% CI)	Not estimable
27.2 Wavelength 820, 830 nm	0	0	Mean Difference (IV, Fixed, 95% CI)	Not estimable
27.3 Wavelength 650-950 nm	1	35	Mean Difference (IV, Fixed, 95% CI)	20.35 [-64.36, 105.06]

Comparison 9. Subgroup Dosage Analysis

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Pain VAS ≤ 3 J/cm ²	2	94	Mean Difference (IV, Fixed, 95% CI)	-1.22 [-1.81, -0.63]
2 Pain VAS > 3 J/cm ²	2	75	Mean Difference (IV, Fixed, 95% CI)	-0.47 [-1.59, 0.65]
3 ESR low dose	1	22	Mean Difference (IV, Fixed, 95% CI)	-20.01 [-28.69, -11.31]
4 ESR high dose	2	70	Mean Difference (IV, Fixed, 95% CI)	-5.34 [-11.35, 0.68]
5 CRP low dose	1	22	Mean Difference (IV, Fixed, 95% CI)	-120.01 [-190.13, -49.87]
6 CRP high dose	1	35	Mean Difference (IV, Fixed, 95% CI)	9.88 [-7.27, 27.03]

7 HAQ high dose	2	75	Mean Difference (IV, Fixed, 95% CI)	0.93 [-2.01, 3.87]
8 Hemoglobin high dose	2	70	Mean Difference (IV, Fixed, 95% CI)	0.47 [0.01, 0.93]
9 Morning stiffness (hours) high dose	3	110	Mean Difference (IV, Fixed, 95% CI)	-27.45 [-51.95, -2.95]
10 Grip strength (kg) low	1	22	Mean Difference (IV, Fixed, 95% CI)	0.06 [-0.49, 1.49]
11 Grip strength (kg) high	2	75	Mean Difference (IV, Fixed, 95% CI)	7.71 [0.15, 15.27]
12 MCP swelling (cm) low	2	75	Mean Difference (IV, Fixed, 95% CI)	0.53 [-1.12, 2.18]
13 Flexibility tip to palm (cm) low	1	22	Mean Difference (IV, Fixed, 95% CI)	-1.26 [-1.68, -0.82]
14 Flexibility tip to palm high dose	1	35	Mean Difference (IV, Fixed, 95% CI)	-6.01 [-11.25, -0.75]

Comparison 10. Laser vs Placebo- contralateral control (End of treatment)


Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Number of patients improved in pain relief	1	34	Risk Ratio (M-H, Fixed, 95% CI)	13.01 [0.79, 214.05]
2 Number of patients improved in morning stiffness	1	34	Risk Ratio (M-H, Fixed, 95% CI)	4.01 [0.50, 32.20]

Analysis 1.1. Comparison 1 Laser vs Placebo- End of treatment (approx 10 wks), Outcome 1 Pain (Visual Analogue Scale 10 cm).

Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 1 Laser vs Placebo- End of treatment (approx 10 wks)

Outcome: 1 Pain (Visual Analogue Scale 10 cm)




Analysis 1.2. Comparison 1 Laser vs Placebo- End of treatment (approx 10 wks), Outcome 2 Pain (0-12 scale).

Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 1 Laser vs Placebo- End of treatment (approx 10 wks)

Outcome: 2 Pain (0-12 scale)




Analysis 1.3. Comparison 1 Laser vs Placebo- End of treatment (approx 10 wks), Outcome 3 McGill Pain questionnaire.

Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 1 Laser vs Placebo- End of treatment (approx 10 wks)

Outcome: 3 McGill Pain questionnaire




Analysis 1.4. Comparison 1 Laser vs Placebo- End of treatment (approx 10 wks), Outcome 4 Ritchie Index.

Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 1 Laser vs Placebo- End of treatment (approx 10 wks)

Outcome: 4 Ritchie Index




Analysis 1.5. Comparison 1 Laser vs Placebo- End of treatment (approx 10 wks), Outcome 5 Health Assessment Questionnaire (HAQ).

Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 1 Laser vs Placebo- End of treatment (approx 10 wks)

Outcome: 5 Health Assessment Questionnaire (HAQ)




Analysis I.6. Comparison I Laser vs Placebo- End of treatment (approx 10 wks), Outcome 6 MCP ROM (degrees).

Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: I Laser vs Placebo- End of treatment (approx 10 wks)

Outcome: 6 MCP ROM (degrees)




Analysis I.7. Comparison I Laser vs Placebo- End of treatment (approx 10 wks), Outcome 7 PIP ROM (degrees).

Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: I Laser vs Placebo- End of treatment (approx 10 wks)

Outcome: 7 PIP ROM (degrees)




Analysis 1.8. Comparison 1 Laser vs Placebo- End of treatment (approx 10 wks), Outcome 8 Knee ROM.

Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 1 Laser vs Placebo- End of treatment (approx 10 wks)


Outcome: 8 Knee ROM



Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 1 Laser vs Placebo- End of treatment (approx 10 wks)


Outcome: 8 Knee ROM



Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: I Laser vs Placebo- End of treatment (approx 10 wks)

Outcome: 8 Knee ROM




Analysis I.9. Comparison I Laser vs Placebo- End of treatment (approx 10 wks), Outcome 9 Ankle ROM.

Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: I Laser vs Placebo- End of treatment (approx 10 wks)


Outcome: 9 Ankle ROM



Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: I Laser vs Placebo- End of treatment (approx 10 wks)


Outcome: 9 Ankle ROM



Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: I Laser vs Placebo- End of treatment (approx 10 wks)

Outcome: 9 Ankle ROM




Analysis I.10. Comparison I Laser vs Placebo- End of treatment (approx 10 wks), Outcome 10 Flexibility- tip to palm distance (cm).

Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: I Laser vs Placebo- End of treatment (approx 10 wks)

Outcome: 10 Flexibility- tip to palm distance (cm)




Analysis I.11. Comparison I Laser vs Placebo- End of treatment (approx 10 wks), Outcome 11 Morning stiffness duration (min).

Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: I Laser vs Placebo- End of treatment (approx 10 wks)

Outcome: 11 Morning stiffness duration (min)




Analysis 1.12. Comparison 1 Laser vs Placebo- End of treatment (approx 10 wks), Outcome 12 Morning stiffness not improved.

Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 1 Laser vs Placebo- End of treatment (approx 10 wks)

Outcome: 12 Morning stiffness not improved




Analysis 1.13. Comparison 1 Laser vs Placebo- End of treatment (approx 10 wks), Outcome 13 Rheumatoid factor positive.

Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 1 Laser vs Placebo- End of treatment (approx 10 wks)

Outcome: 13 Rheumatoid factor positive




Analysis I.14. Comparison I Laser vs Placebo- End of treatment (approx 10 wks), Outcome 14 Grip strength.

Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: I Laser vs Placebo- End of treatment (approx 10 wks)


Outcome: 14 Grip strength



Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: I Laser vs Placebo- End of treatment (approx 10 wks)


Outcome: 14 Grip strength



Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: I Laser vs Placebo- End of treatment (approx 10 wks)

Outcome: 14 Grip strength




Analysis 1.15. Comparison I Laser vs Placebo- End of treatment (approx 10 wks), Outcome 15 Suprapatellar swelling - right knee (cm).

Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: I Laser vs Placebo- End of treatment (approx 10 wks)

Outcome: 15 Suprapatellar swelling - right knee (cm)




Analysis I.16. Comparison I Laser vs Placebo- End of treatment (approx 10 wks), Outcome 16 Suprapatellar swelling- left knee (cm).

Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: I Laser vs Placebo- End of treatment (approx 10 wks)

Outcome: 16 Suprapatellar swelling- left knee (cm)




Analysis I.17. Comparison I Laser vs Placebo- End of treatment (approx 10 wks), Outcome 17 MCP swelling (cm).

Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: I Laser vs Placebo- End of treatment (approx 10 wks)

Outcome: 17 MCP swelling (cm)




Analysis 1.18. Comparison 1 Laser vs Placebo- End of treatment (approx 10 wks), Outcome 18 PIP swelling (cm).

Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 1 Laser vs Placebo- End of treatment (approx 10 wks)

Outcome: 18 PIP swelling (cm)




Analysis 1.19. Comparison 1 Laser vs Placebo- End of treatment (approx 10 wks), Outcome 19 Walking speed (sec).

Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 1 Laser vs Placebo- End of treatment (approx 10 wks)

Outcome: 19 Walking speed (sec)




Analysis 1.20. Comparison I Laser vs Placebo- End of treatment (approx 10 wks), Outcome 20 Fibrinogen.

Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: I Laser vs Placebo- End of treatment (approx 10 wks)

Outcome: 20 Fibrinogen




Analysis 1.21. Comparison I Laser vs Placebo- End of treatment (approx 10 wks), Outcome 21 Leukocytes.

Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: I Laser vs Placebo- End of treatment (approx 10 wks)

Outcome: 21 Leukocytes




Analysis I.22. Comparison I Laser vs Placebo- End of treatment (approx 10 wks), Outcome 22 Lymphocytes.

Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: I Laser vs Placebo- End of treatment (approx 10 wks)

Outcome: 22 Lymphocytes




Analysis I.23. Comparison I Laser vs Placebo- End of treatment (approx 10 wks), Outcome 23 ESR (mm/hr).

Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: I Laser vs Placebo- End of treatment (approx 10 wks)

Outcome: 23 ESR (mm/hr)




Analysis 1.24. Comparison I Laser vs Placebo- End of treatment (approx 10 wks), Outcome 24 CRP (g/mL).

Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: I Laser vs Placebo- End of treatment (approx 10 wks)

Outcome: 24 CRP (g/mL)




Analysis 1.25. Comparison I Laser vs Placebo- End of treatment (approx 10 wks), Outcome 25 Hemoglobin.

Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: I Laser vs Placebo- End of treatment (approx 10 wks)

Outcome: 25 Hemoglobin




Analysis 1.26. Comparison 1 Laser vs Placebo- End of treatment (approx 10 wks), Outcome 26 Platelets.

Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 1 Laser vs Placebo- End of treatment (approx 10 wks)

Outcome: 26 Platelets




Analysis 2.1. Comparison 2 Laser vs Placebo- End of follow-up (approx 20 wks), Outcome 1 Pain.

Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 2 Laser vs Placebo- End of follow-up (approx 20 wks)


Outcome: 1 Pain



Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 2 Laser vs Placebo- End of follow-up (approx 20 wks)

Outcome: 1 Pain




Analysis 2.2. Comparison 2 Laser vs Placebo- End of follow-up (approx 20 wks), Outcome 2 McGill Pain questionnaire.

Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 2 Laser vs Placebo- End of follow-up (approx 20 wks)

Outcome: 2 McGill Pain questionnaire




Analysis 2.3. Comparison 2 Laser vs Placebo- End of follow-up (approx 20 wks), Outcome 3 Health Assessment Questionnaire (HAQ).

Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 2 Laser vs Placebo- End of follow-up (approx 20 wks)

Outcome: 3 Health Assessment Questionnaire (HAQ)




Analysis 2.4. Comparison 2 Laser vs Placebo- End of follow-up (approx 20 wks), Outcome 4 Ritchie Index.

Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 2 Laser vs Placebo- End of follow-up (approx 20 wks)

Outcome: 4 Ritchie Index




Analysis 2.5. Comparison 2 Laser vs Placebo- End of follow-up (approx 20 wks), Outcome 5 PIP ROM (degrees).

Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 2 Laser vs Placebo- End of follow-up (approx 20 wks)

Outcome: 5 PIP ROM (degrees)




Analysis 2.6. Comparison 2 Laser vs Placebo- End of follow-up (approx 20 wks), Outcome 6 Knee ROM.

Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 2 Laser vs Placebo- End of follow-up (approx 20 wks)


Outcome: 6 Knee ROM



Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 2 Laser vs Placebo- End of follow-up (approx 20 wks)


Outcome: 6 Knee ROM



Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 2 Laser vs Placebo- End of follow-up (approx 20 wks)

Outcome: 6 Knee ROM




Analysis 2.7. Comparison 2 Laser vs Placebo- End of follow-up (approx 20 wks), Outcome 7 Ankle ROM.

Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 2 Laser vs Placebo- End of follow-up (approx 20 wks)


Outcome: 7 Ankle ROM



Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 2 Laser vs Placebo- End of follow-up (approx 20 wks)


Outcome: 7 Ankle ROM



Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 2 Laser vs Placebo- End of follow-up (approx 20 wks)

Outcome: 7 Ankle ROM




Analysis 2.8. Comparison 2 Laser vs Placebo- End of follow-up (approx 20 wks), Outcome 8 MCP ROM (degrees).

Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 2 Laser vs Placebo- End of follow-up (approx 20 wks)

Outcome: 8 MCP ROM (degrees)




Analysis 2.9. Comparison 2 Laser vs Placebo- End of follow-up (approx 20 wks), Outcome 9 Morning stiffness duration (min).

Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 2 Laser vs Placebo- End of follow-up (approx 20 wks)

Outcome: 9 Morning stiffness duration (min)




Analysis 2.10. Comparison 2 Laser vs Placebo- End of follow-up (approx 20 wks), Outcome 10 Walking speed (sec).

Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 2 Laser vs Placebo- End of follow-up (approx 20 wks)

Outcome: 10 Walking speed (sec)




Analysis 2.11. Comparison 2 Laser vs Placebo- End of follow-up (approx 20 wks), Outcome 11 Grip strength.

Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 2 Laser vs Placebo- End of follow-up (approx 20 wks)


Outcome: 11 Grip strength



Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 2 Laser vs Placebo- End of follow-up (approx 20 wks)

Outcome: 11 Grip strength




Analysis 2.12. Comparison 2 Laser vs Placebo- End of follow-up (approx 20 wks), Outcome 12 Suprapatellar swelling- left knee (cm).

Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 2 Laser vs Placebo- End of follow-up (approx 20 wks)

Outcome: 12 Suprapatellar swelling- left knee (cm)




Analysis 2.13. Comparison 2 Laser vs Placebo- End of follow-up (approx 20 wks), Outcome 13 Suprapatellar swelling - right knee (cm).

Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 2 Laser vs Placebo- End of follow-up (approx 20 wks)

Outcome: 13 Suprapatellar swelling - right knee (cm)




Analysis 2.14. Comparison 2 Laser vs Placebo- End of follow-up (approx 20 wks), Outcome 14 MCP swelling (cm).

Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 2 Laser vs Placebo- End of follow-up (approx 20 wks)

Outcome: 14 MCP swelling (cm)




Analysis 2.15. Comparison 2 Laser vs Placebo- End of follow-up (approx 20 wks), Outcome 15 PIP swelling (cm).

Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 2 Laser vs Placebo- End of follow-up (approx 20 wks)

Outcome: 15 PIP swelling (cm)




Analysis 2.16. Comparison 2 Laser vs Placebo- End of follow-up (approx 20 wks), Outcome 16 Thermographic index.

Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 2 Laser vs Placebo- End of follow-up (approx 20 wks)

Outcome: 16 Thermographic index




Analysis 2.17. Comparison 2 Laser vs Placebo- End of follow-up (approx 20 wks), Outcome 17 Rheumatoid factor positive.

Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 2 Laser vs Placebo- End of follow-up (approx 20 wks)

Outcome: 17 Rheumatoid factor positive




Analysis 2.18. Comparison 2 Laser vs Placebo- End of follow-up (approx 20 wks), Outcome 18 ESR (mm/hr).

Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 2 Laser vs Placebo- End of follow-up (approx 20 wks)

Outcome: 18 ESR (mm/hr)




Analysis 2.19. Comparison 2 Laser vs Placebo- End of follow-up (approx 20 wks), Outcome 19 CRP (g/mL).

Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 2 Laser vs Placebo- End of follow-up (approx 20 wks)

Outcome: 19 CRP (g/mL)




Analysis 2.20. Comparison 2 Laser vs Placebo- End of follow-up (approx 20 wks), Outcome 20 Hemoglobin.

Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 2 Laser vs Placebo- End of follow-up (approx 20 wks)

Outcome: 20 Hemoglobin




Analysis 2.21. Comparison 2 Laser vs Placebo- End of follow-up (approx 20 wks), Outcome 21 Platelets.

Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 2 Laser vs Placebo- End of follow-up (approx 20 wks)

Outcome: 21 Platelets




Analysis 3.1. Comparison 3 Treatment length- subgroup analysis for end of treatment results, Outcome 1 Pain (10 cm VAS or 0-12 scale).

Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 3 Treatment length- subgroup analysis for end of treatment results


Outcome: 1 Pain (10 cm VAS or 0-12 scale)



Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 3 Treatment length- subgroup analysis for end of treatment results


Outcome: 1 Pain (10 cm VAS or 0-12 scale)



Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 3 Treatment length- subgroup analysis for end of treatment results

Outcome: 1 Pain (10 cm VAS or 0-12 scale)




Analysis 4.1. Comparison 4 Treatment length- subgroup analysis for end of follow up results, Outcome 1 Pain (10 cm VAS or 0-12 scale).

Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 4 Treatment length- subgroup analysis for end of follow up results


Outcome: 1 Pain (10 cm VAS or 0-12 scale)



Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 4 Treatment length- subgroup analysis for end of follow up results

Outcome: 1 Pain (10 cm VAS or 0-12 scale)




Analysis 5.1. Comparison 5 Methodologic quality: low (<3) vs high (=> 3), Outcome 1 Pain at end of treatment.


Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 5 Methodologic quality: low (<3) vs high (=> 3)


Outcome: 1 Pain at end of treatment



(Continued . . .)




Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis
 Comparison: 5 Methodologic quality: low (<3) vs high (=> 3)
 Outcome: 1 Pain at end of treatment



Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 5 Methodologic quality: low (<3) vs high (=> 3)

Outcome: 1 Pain at end of treatment




Analysis 6.1. Comparison 6 Laser vs Placebo - Joint vs Nerve - end of treatment, Outcome 1 Pain (10 cm VAS or 0-12 scale).


Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 6 Laser vs Placebo - Joint vs Nerve - end of treatment

Outcome: 1 Pain (10 cm VAS or 0-12 scale)




(Continued ...)



Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 6 Laser vs Placebo - Joint vs Nerve - end of treatment


Outcome: 1 Pain (10 cm VAS or 0-12 scale)



Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 6 Laser vs Placebo - Joint vs Nerve - end of treatment

Outcome: 1 Pain (10 cm VAS or 0-12 scale)




Analysis 7.1. Comparison 7 Laser vs Placebo - Joint vs Nerve - end of follow up, Outcome 1 Pain (10 cm VAS or 0-12 scale).

Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 7 Laser vs Placebo - Joint vs Nerve - end of follow up


Outcome: 1 Pain (10 cm VAS or 0-12 scale)



Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 7 Laser vs Placebo - Joint vs Nerve - end of follow up

Outcome: 1 Pain (10 cm VAS or 0-12 scale)




Analysis 8.1. Comparison 8 Laser vs Placebo- Wavelength analysis - End of treatment, Outcome 1 Pain.

Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 8 Laser vs Placebo- Wavelength analysis - End of treatment


Outcome: 1 Pain



Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 8 Laser vs Placebo- Wavelength analysis - End of treatment


Outcome: 1 Pain



Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 8 Laser vs Placebo- Wavelength analysis - End of treatment


Outcome: 1 Pain



Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 8 Laser vs Placebo- Wavelength analysis - End of treatment

Outcome: 1 Pain




Analysis 8.2. Comparison 8 Laser vs Placebo- Wavelength analysis - End of treatment, Outcome 2 McGill Pain questionnaire.

Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 8 Laser vs Placebo- Wavelength analysis - End of treatment


Outcome: 2 McGill Pain questionnaire



Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 8 Laser vs Placebo- Wavelength analysis - End of treatment

Outcome: 2 McGill Pain questionnaire




Analysis 8.3. Comparison 8 Laser vs Placebo- Wavelength analysis - End of treatment, Outcome 3 Ritchie Index.

Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 8 Laser vs Placebo- Wavelength analysis - End of treatment


Outcome: 3 Ritchie Index



Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 8 Laser vs Placebo- Wavelength analysis - End of treatment

Outcome: 3 Ritchie Index




Analysis 8.4. Comparison 8 Laser vs Placebo- Wavelength analysis - End of treatment, Outcome 4 Health Assessment Questionnaire (HAQ).

Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 8 Laser vs Placebo- Wavelength analysis - End of treatment


Outcome: 4 Health Assessment Questionnaire (HAQ)



Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 8 Laser vs Placebo- Wavelength analysis - End of treatment


Outcome: 4 Health Assessment Questionnaire (HAQ)



Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 8 Laser vs Placebo- Wavelength analysis - End of treatment

Outcome: 4 Health Assessment Questionnaire (HAQ)




Analysis 8.5. Comparison 8 Laser vs Placebo- Wavelength analysis - End of treatment, Outcome 5 MCP ROM (degrees).

Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 8 Laser vs Placebo- Wavelength analysis - End of treatment


Outcome: 5 MCP ROM (degrees)



Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 8 Laser vs Placebo- Wavelength analysis - End of treatment

Outcome: 5 MCP ROM (degrees)




Analysis 8.6. Comparison 8 Laser vs Placebo- Wavelength analysis - End of treatment, Outcome 6 PIP ROM (degrees).

Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 8 Laser vs Placebo- Wavelength analysis - End of treatment


Outcome: 6 PIP ROM (degrees)



Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 8 Laser vs Placebo- Wavelength analysis - End of treatment

Outcome: 6 PIP ROM (degrees)




Analysis 8.7. Comparison 8 Laser vs Placebo- Wavelength analysis - End of treatment, Outcome 7 Left knee Range of motion.

Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 8 Laser vs Placebo- Wavelength analysis - End of treatment


Outcome: 7 Left knee Range of motion



Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 8 Laser vs Placebo- Wavelength analysis - End of treatment

Outcome: 7 Left knee Range of motion




Analysis 8.8. Comparison 8 Laser vs Placebo- Wavelength analysis - End of treatment, Outcome 8 Right knee ROM.

Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 8 Laser vs Placebo- Wavelength analysis - End of treatment


Outcome: 8 Right knee ROM



Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 8 Laser vs Placebo- Wavelength analysis - End of treatment

Outcome: 8 Right knee ROM




Analysis 8.9. Comparison 8 Laser vs Placebo- Wavelength analysis - End of treatment, Outcome 9 Left ankle ROM.

Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 8 Laser vs Placebo- Wavelength analysis - End of treatment


Outcome: 9 Left ankle ROM



Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 8 Laser vs Placebo- Wavelength analysis - End of treatment

Outcome: 9 Left ankle ROM




Analysis 8.10. Comparison 8 Laser vs Placebo- Wavelength analysis - End of treatment, Outcome 10 Right ankle ROM.

Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 8 Laser vs Placebo- Wavelength analysis - End of treatment


Outcome: 10 Right ankle ROM



Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 8 Laser vs Placebo- Wavelength analysis - End of treatment

Outcome: 10 Right ankle ROM




Analysis 8.11. Comparison 8 Laser vs Placebo- Wavelength analysis - End of treatment, Outcome 11 Flexibility- tip to palm distance (cm).

Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 8 Laser vs Placebo- Wavelength analysis - End of treatment


Outcome: 11 Flexibility- tip to palm distance (cm)



Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 8 Laser vs Placebo- Wavelength analysis - End of treatment

Outcome: 11 Flexibility- tip to palm distance (cm)




Analysis 8.12. Comparison 8 Laser vs Placebo- Wavelength analysis - End of treatment, Outcome 12 Morning stiffness duration (min).

Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 8 Laser vs Placebo- Wavelength analysis - End of treatment


Outcome: 12 Morning stiffness duration (min)



Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 8 Laser vs Placebo- Wavelength analysis - End of treatment


Outcome: 12 Morning stiffness duration (min)



Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 8 Laser vs Placebo- Wavelength analysis - End of treatment

Outcome: 12 Morning stiffness duration (min)




Analysis 8.13. Comparison 8 Laser vs Placebo- Wavelength analysis - End of treatment, Outcome 13 Morning stiffness not improved.

Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 8 Laser vs Placebo- Wavelength analysis - End of treatment


Outcome: 13 Morning stiffness not improved



Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 8 Laser vs Placebo- Wavelength analysis - End of treatment

Outcome: 13 Morning stiffness not improved




Analysis 8.14. Comparison 8 Laser vs Placebo- Wavelength analysis - End of treatment, Outcome 14 Rheumatoid factor positive.

Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 8 Laser vs Placebo- Wavelength analysis - End of treatment


Outcome: 14 Rheumatoid factor positive



Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 8 Laser vs Placebo- Wavelength analysis - End of treatment

Outcome: 14 Rheumatoid factor positive




Analysis 8.15. Comparison 8 Laser vs Placebo- Wavelength analysis - End of treatment, Outcome 15 Grip strength (mmHg, KPa, kg).

Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 8 Laser vs Placebo- Wavelength analysis - End of treatment


Outcome: 15 Grip strength (mmHg, KPa, kg)



Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 8 Laser vs Placebo- Wavelength analysis - End of treatment

Outcome: 15 Grip strength (mmHg, KPa, kg)




Analysis 8.16. Comparison 8 Laser vs Placebo- Wavelength analysis - End of treatment, Outcome 16 Suprapatellar swelling - right knee (cm).

Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 8 Laser vs Placebo- Wavelength analysis - End of treatment


Outcome: 16 Suprapatellar swelling - right knee (cm)



Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 8 Laser vs Placebo- Wavelength analysis - End of treatment

Outcome: 16 Suprapatellar swelling - right knee (cm)




Analysis 8.17. Comparison 8 Laser vs Placebo- Wavelength analysis - End of treatment, Outcome 17 Suprapatellar swelling- left knee (cm).

Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 8 Laser vs Placebo- Wavelength analysis - End of treatment


Outcome: 17 Suprapatellar swelling- left knee (cm)



Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 8 Laser vs Placebo- Wavelength analysis - End of treatment

Outcome: 17 Suprapatellar swelling- left knee (cm)




Analysis 8.18. Comparison 8 Laser vs Placebo- Wavelength analysis - End of treatment, Outcome 18 MCP swelling (cm).

Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 8 Laser vs Placebo- Wavelength analysis - End of treatment


Outcome: 18 MCP swelling (cm)



Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 8 Laser vs Placebo- Wavelength analysis - End of treatment

Outcome: 18 MCP swelling (cm)




Analysis 8.19. Comparison 8 Laser vs Placebo- Wavelength analysis - End of treatment, Outcome 19 PIP swelling (cm).

Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 8 Laser vs Placebo- Wavelength analysis - End of treatment


Outcome: 19 PIP swelling (cm)



Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 8 Laser vs Placebo- Wavelength analysis - End of treatment

Outcome: 19 PIP swelling (cm)




Analysis 8.20. Comparison 8 Laser vs Placebo- Wavelength analysis - End of treatment, Outcome 20 Walking speed (sec).

Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 8 Laser vs Placebo- Wavelength analysis - End of treatment


Outcome: 20 Walking speed (sec)



Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 8 Laser vs Placebo- Wavelength analysis - End of treatment

Outcome: 20 Walking speed (sec)




Analysis 8.21. Comparison 8 Laser vs Placebo- Wavelength analysis - End of treatment, Outcome 21 Fibrinogen.

Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 8 Laser vs Placebo- Wavelength analysis - End of treatment


Outcome: 21 Fibrinogen



Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 8 Laser vs Placebo- Wavelength analysis - End of treatment

Outcome: 21 Fibrinogen




Analysis 8.22. Comparison 8 Laser vs Placebo- Wavelength analysis - End of treatment, Outcome 22 Leukocytes.

Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 8 Laser vs Placebo- Wavelength analysis - End of treatment


Outcome: 22 Leukocytes



Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 8 Laser vs Placebo- Wavelength analysis - End of treatment

Outcome: 22 Leukocytes




Analysis 8.23. Comparison 8 Laser vs Placebo- Wavelength analysis - End of treatment, Outcome 23 Lymphocytes.

Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 8 Laser vs Placebo- Wavelength analysis - End of treatment


Outcome: 23 Lymphocytes



Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 8 Laser vs Placebo- Wavelength analysis - End of treatment

Outcome: 23 Lymphocytes




Analysis 8.24. Comparison 8 Laser vs Placebo- Wavelength analysis - End of treatment, Outcome 24 ESR (mm/hr).

Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 8 Laser vs Placebo- Wavelength analysis - End of treatment


Outcome: 24 ESR (mm/hr)



Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 8 Laser vs Placebo- Wavelength analysis - End of treatment


Outcome: 24 ESR (mm/hr)



Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 8 Laser vs Placebo- Wavelength analysis - End of treatment

Outcome: 24 ESR (mm/hr)




Analysis 8.25. Comparison 8 Laser vs Placebo- Wavelength analysis - End of treatment, Outcome 25 CRP (g/mL).

Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis


Comparison: 8 Laser vs Placebo- Wavelength analysis - End of treatment

Outcome: 25 CRP (g/mL)



(Continued ...)


(... Continued)



Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 8 Laser vs Placebo- Wavelength analysis - End of treatment


Outcome: 25 CRP (g/mL)



Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 8 Laser vs Placebo- Wavelength analysis - End of treatment

Outcome: 25 CRP (g/mL)




Analysis 8.26. Comparison 8 Laser vs Placebo- Wavelength analysis - End of treatment, Outcome 26 Hemoglobin.

Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 8 Laser vs Placebo- Wavelength analysis - End of treatment


Outcome: 26 Hemoglobin



Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 8 Laser vs Placebo- Wavelength analysis - End of treatment


Outcome: 26 Hemoglobin



Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 8 Laser vs Placebo- Wavelength analysis - End of treatment

Outcome: 26 Hemoglobin




Analysis 8.27. Comparison 8 Laser vs Placebo- Wavelength analysis - End of treatment, Outcome 27 Platelets.

Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 8 Laser vs Placebo- Wavelength analysis - End of treatment


Outcome: 27 Platelets



Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 8 Laser vs Placebo- Wavelength analysis - End of treatment

Outcome: 27 Platelets




Analysis 9.1. Comparison 9 Subgroup Dosage Analysis, Outcome 1 Pain VAS <= 3 J/cm2.

Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 9 Subgroup Dosage Analysis

Outcome: 1 Pain VAS <= 3 J/cm2




Analysis 9.2. Comparison 9 Subgroup Dosage Analysis, Outcome 2 Pain VAS > 3 J/cm2.

Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 9 Subgroup Dosage Analysis

Outcome: 2 Pain VAS > 3 J/cm2




Analysis 9.3. Comparison 9 Subgroup Dosage Analysis, Outcome 3 ESR low dose.

Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 9 Subgroup Dosage Analysis

Outcome: 3 ESR low dose




Analysis 9.4. Comparison 9 Subgroup Dosage Analysis, Outcome 4 ESR high dose.

Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 9 Subgroup Dosage Analysis

Outcome: 4 ESR high dose




Analysis 9.5. Comparison 9 Subgroup Dosage Analysis, Outcome 5 CRP low dose.

Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 9 Subgroup Dosage Analysis

Outcome: 5 CRP low dose




Analysis 9.6. Comparison 9 Subgroup Dosage Analysis, Outcome 6 CRP high dose.

Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 9 Subgroup Dosage Analysis

Outcome: 6 CRP high dose




Analysis 9.7. Comparison 9 Subgroup Dosage Analysis, Outcome 7 HAQ high dose.

Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 9 Subgroup Dosage Analysis

Outcome: 7 HAQ high dose




Analysis 9.8. Comparison 9 Subgroup Dosage Analysis, Outcome 8 Hemoglobin high dose.

Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 9 Subgroup Dosage Analysis

Outcome: 8 Hemoglobin high dose




Analysis 9.9. Comparison 9 Subgroup Dosage Analysis, Outcome 9 Morning stiffness (hours) high dose.

Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 9 Subgroup Dosage Analysis

Outcome: 9 Morning stiffness (hours) high dose




Analysis 9.10. Comparison 9 Subgroup Dosage Analysis, Outcome 10 Grip strength (kg) low.

Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 9 Subgroup Dosage Analysis

Outcome: 10 Grip strength (kg) low




Analysis 9.11. Comparison 9 Subgroup Dosage Analysis, Outcome 11 Grip strength (kg) high.

Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 9 Subgroup Dosage Analysis

Outcome: 11 Grip strength (kg) high




Analysis 9.12. Comparison 9 Subgroup Dosage Analysis, Outcome 12 MCP swelling (cm) low.

Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 9 Subgroup Dosage Analysis

Outcome: 12 MCP swelling (cm) low




Analysis 9.13. Comparison 9 Subgroup Dosage Analysis, Outcome 13 Flexibility tip to palm (cm) low.

Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 9 Subgroup Dosage Analysis

Outcome: 13 Flexibility tip to palm (cm) low




Analysis 9.14. Comparison 9 Subgroup Dosage Analysis, Outcome 14 Flexibility tip to palm high dose.

Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 9 Subgroup Dosage Analysis

Outcome: 14 Flexibility tip to palm high dose




**Analysis 10.1. Comparison 10 Laser vs Placebo- contralateral control (End of treatment), Outcome 1
Number of patients improved in pain relief.**

Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 10 Laser vs Placebo- contralateral control (End of treatment)

Outcome: 1 Number of patients improved in pain relief




**Analysis 10.2. Comparison 10 Laser vs Placebo- contralateral control (End of treatment), Outcome 2
Number of patients improved in morning stiffness.**

Review: Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis

Comparison: 10 Laser vs Placebo- contralateral control (End of treatment)

Outcome: 2 Number of patients improved in morning stiffness



APPENDICES

Appendix I. Full search strategy

- 1 degenerative arthritis.tw.
- 2 exp arthritis, rheumatoid/
- 3 rheumatoid arthritis.tw.
- 4 rheumatism.tw.
- 5 arthritis, juvenile rheumatoid/
- 6 caplan's syndrome.tw.
- 7 felty's syndrome.tw.
- 8 rheumatoid.tw.
- 9 ankylosing spondylitis.tw.
- 10 arthrosis.tw.
- 11 sjogren\$.tw.
- 12 or/1-11
- 13 laser\$.sh.
- 14 laser\$.tw.
- 15 exp light/
- 16 infrared.tw.
- 17 ultraviolet.tw.
- 18 monochromatic.tw.
- 19 or/13-18
- 20 12 and 19
- 21 clinical trial.pt.
- 22 randomized controlled trial.pt.
- 23 tu.fs.
- 24 dt.fs.
- 25 random\$.tw.
- 26 placebo\$.tw.
- 27 ((sing\$ or doubl\$ or tripl\$) adj (masked or blind\$)).tw
- 28 sham.tw.
- 29 or/21-28
- 30 20 and 29

WHAT'S NEW

Last assessed as up-to-date: 21 August 2005.

29 September 2008	Amended	Converted to new review format. CMSG ID: C049-R
-------------------	---------	--

HISTORY

Review first published: Issue 4, 1998

CONTRIBUTIONS OF AUTHORS

Lucie Brosseau is responsible for the content of the paper and revisions.

Vivian Robinson conducted data extraction, analyses and contributed to the manuscript.

George Wells helped interpret the data from the individual trials and the statistically pooled results.

Arne Gam and Rob de Bie contributed to the literature selection and data extraction.

Katherine Harman provided content expertise in pain control theory.

Michelle Morin contributed to the theoretical framework that was used for data extraction.

Bev Shea helped with quality assessment and editing of the manuscript.

Peter Tugwell contributed to the clinical interpretation and presentation of results.

DECLARATIONS OF INTEREST

None known

SOURCES OF SUPPORT

Internal sources

- University of Ottawa, Canada.

External sources

- No sources of support supplied

INDEX TERMS

Medical Subject Headings (MeSH)

Arthritis, Rheumatoid [*radiotherapy]; *Laser Therapy, Low-Level; Randomized Controlled Trials as Topic

MeSH check words

Humans